

Stable Directions for Degenerate Excited States of Nonlinear Schrödinger Equations

Stephen Gustafson, Tuoc Van Phan

April 13, 2010

Abstract

We consider nonlinear Schrödinger equations, $i\partial_t\psi = H_0\psi + \lambda|\psi|^2\psi$ in $\mathbb{R}^3 \times [0, \infty)$, where $H_0 = -\Delta + V$, $\lambda = \pm 1$, the potential V is radial and spatially decaying, and the linear Hamiltonian H_0 has only two eigenvalues $e_0 < e_1 < 0$, where e_0 is simple, and e_1 has multiplicity three. We show that there exist two branches of small “nonlinear excited state” standing-wave solutions, and in both the resonant ($e_0 < 2e_1$) and non-resonant ($e_0 > 2e_1$) cases, we construct certain finite-codimension regions of the phase space consisting of solutions converging to these excited states at time infinity (“stable directions”).

Key words. Asymptotic dynamics, Nonlinear excited states, Schrödinger equations.

AMS subject classifications. 35Q40; 35Q55.

1 Introduction

We consider the nonlinear Schrödinger equation

$$i\partial_t\psi = H_0\psi + \lambda|\psi|^2\psi \tag{1.1}$$

which arises in several physical settings, including many-body quantum systems, and optics. Here the wave function $\psi = \psi(x, t)$ is complex-valued,

$$\psi : \mathbb{R}^3 \times [0, \infty) \rightarrow \mathbb{C},$$

and the linear Hamiltonian is

$$H_0 := -\Delta + V(x),$$

where $V : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a smooth, spatially decaying potential function. We take $\lambda = \pm 1$. Equation (1.1) can be expressed as a Hamiltonian system

$$i\partial_t \psi = \frac{\partial \mathcal{E}[\psi, \bar{\psi}]}{\partial \bar{\psi}},$$

where the Hamiltonian energy is defined as

$$\mathcal{E}[\psi] = \mathcal{E}[\psi, \bar{\psi}] = \int \left(\frac{1}{2} \nabla \psi \cdot \nabla \bar{\psi} + \frac{1}{2} V(x) \psi \bar{\psi} + \frac{\lambda}{4} \psi^2 \bar{\psi}^2 \right) dx.$$

Since the energy is invariant under the time-translation $t \mapsto t + t_0$ ($t_0 \in \mathbb{R}$) and the phase rotation

$$\psi \mapsto e^{ir} \psi, \quad r \in \mathbb{R},$$

the energy and the particle number

$$\mathcal{N}[\psi] := \int |\psi(x)|^2 dx$$

are constant in time for any smooth solution $\psi(t) = \psi(., t) \in H^1(\mathbb{R}^3)$ of (1.1):

$$\mathcal{E}[\psi(t)] = \mathcal{E}[\psi(0)], \quad \mathcal{N}[\psi(t)] = \mathcal{N}[\psi(0)]$$

The global well-posedness of solutions with $\|\psi(0)\|_{H^1}$ small can be established easily by using these conserved quantities, and a continuity argument.

A very important feature of the equation (1.1) is that it can have localized “standing wave” or “nonlinear bound state” solutions. These are solutions of the form $\psi(x, t) = e^{-iEt} Q(x)$, where the profile function Q must therefore solve the equation

$$H_0 Q + \lambda |Q|^2 Q = EQ. \tag{1.2}$$

The solutions of (1.2) are critical points of the Hamiltonian $\mathcal{E}$ subject to the constraint that the L^2 -norm of Q is fixed. We may obtain *small* solutions of (1.2) as bifurcations along the discrete eigenvalues of H_0 . In this paper, we assume that H_0 has two discrete eigenvalues $e_0 < e_1 < 0$. The small solutions with E close to e_0 are called *nonlinear ground states*, while those with E close to e_1 are naturally called *nonlinear excited states*.

Along a simple eigenvalue, the bifurcation problem for finding the corresponding nonlinear solutions (eg. nonlinear ground states) is quite standard, regardless of the multiplicities of the other eigenvalues, see [18, 21, 12]. Along a degenerate eigenvalue (such as e_1 in our setting), however, the problem becomes more delicate due to the interaction between the different directions in the e_1 -eigenspace. Hence our first goal in this paper is to find all of the nonlinear excited states when e_1 is degenerate.

We consider here a *radial* potential V , and we assume that e_1 has multiplicity three, corresponding to the first non-zero angular momentum spherical harmonics:

Assumption A1: *The potential $V(x)$ is spherically symmetric, and the linear Hamiltonian $H_0 = -\Delta + V$ has a discrete eigenvalue $e_1 < 0$ of multiplicity three, with eigenspace*

$$\mathbf{V} := \text{Null}(H_0 - e_1) = \text{span} \{ \phi_1, \phi_2, \phi_3 \}, \quad \phi_j(x) = x_j \varphi(|x|)$$

for a real function φ .

Notice that such a multiplicity-three eigenvalue is a generic possibility in three dimensions with a radial potential.

We will denote the orthogonal projection onto this excited-state eigenspace by

$$\mathbf{P}_1 = \mathbf{P}_1(H_0) := \text{orthogonal projection onto } \mathbf{V}.$$

Since the potential V is radial, equation (1.2) is invariant not only under phase rotation, time translation, and complex conjugation, but also under all spatial rotations and reflections:

$$Q(x) \mapsto Q(\Gamma x), \quad \Gamma \in O(3).$$

This rich structure plays a central role in understanding the existence of nonlinear excited states, as well as the nearby dynamics.

In Section 3, we shall see that there is a *symmetry breaking bifurcation phenomenon* in the bifurcation equations of the nonlinear excited states Q . Here, we shall briefly describe the phenomenon (for the systematic theory, see [10, 11, 17]). By integrating (1.2) against a linear excited state ϕ_j , it follows that there are non-trivial solutions close to $\mathbf{V}$ only if $\lambda(E - e_1) > 0$. We write $Q = \epsilon(v + h)$ for $v \in \mathbf{V}$, $h \in \mathbf{V}^\perp$ and $\epsilon^2 = [E - e_1]/\lambda$. Then, the equation (1.2) is equivalent to

$$\begin{cases} h = \lambda \epsilon^2 [H_0 - e_1]^{-1} (1 - \mathbf{P}_1(H_0)) \{ h - |v + h|^2 (v + h) \} \\ \mathbf{P}_1(H_0) [v - |v + h|^2 (v + h)] = 0 \end{cases}. \quad (1.3)$$

By using the contraction mapping theorem, we obtain the unique small solution $h = h(\epsilon, v)$ of the first equation of (1.3) for sufficiently small $\epsilon > 0$. To solve the bifurcation equation

$$N(\epsilon, v) := \mathbf{P}_1(H_0) [v - |v + h(\epsilon, v)|^2 (v + h(\epsilon, v))] = 0,$$

the standard method is to apply the implicit function theorem. However, due to the invariant structure of N , its derivative $N_v(0, v)$ has a non-trivial kernel

for all $v \in \mathbf{V}$ such that $N(0, v) = 0$. To overcome this, we have to restrict N onto N -invariant subspaces of $\mathbf{V}$, and so eliminate the kernel of the derivative of N . In this way, we obtain two particular representatives, denoted Q_E and $\tilde{Q}_E$, of the family of nonlinear excited states. Then, applying the symmetries – spatial rotation, reflection, phase rotation and complex conjugation – to Q_E and $\tilde{Q}_E$, we obtain all of the nonlinear excited states of (1.2), as summarized in the following theorem, proved in Section 3:

Theorem 1.1 (Existence of degenerate excited states). *There exist two branches of nonlinear excited state solutions to (1.2), with $E - e_1$ small. These branches are generated by applying phase rotations ($Q(x) \mapsto e^{ir}Q(x)$, $r \in \mathbb{R}$) and spatial rotations ($Q(x) \mapsto Q(\Gamma x)$, $\Gamma \in SO(3)$) to two fixed solutions of the forms*

$$Q_E(x) = x_1 f_1(x_1^2, x_2^2 + x_3^2), \quad \tilde{Q}_E(x) = e^{i\theta} f_2(x_1^2 + x_2^2, x_3^2)$$

for real-valued functions f_1, f_2 , and where $\theta \in [0, 2\pi)$ is the angle between the x_1 axis and the vector $x' = (x_1, x_2) \in \mathbb{R}^2$. The solutions Q_E and $\tilde{Q}_E$ lie in the Sobolev space H^2 , and decay exponentially as $|x| \rightarrow \infty$.

Remark 1.1. Another way to state this result: the excited state branches arise as the orbits, under the symmetry group of the equation, of two particular bifurcation curves of solutions. One of these (Q_E) is real, odd in one direction, and invariant under rotations fixing that direction, while the other ($\tilde{Q}_E$) is even in one direction, and “co-rotational” with respect to rotations fixing that direction.

Remark 1.2. The nature of the symmetry group, together with the symmetry properties of Q_E and $\tilde{Q}_E$, imply that each solution branch is a four-dimensional family: one degree of freedom is the eigenvalue E , one is the phase rotation, and two come from spatial rotations. See (3.18) for an explicit expression of the branches in terms of parameters.

Our next goal is to study the dynamics nearby the nonlinear excited states. It is well-known that the family of *ground states* – let us denote them by Q_E^0 , with nonlinear eigenvalue E close to e_0 – is *orbitally stable*; that is the orbit of a ground state under the action of the symmetry group (in this case just phase rotations, since Q_E^0 is radially symmetric) is stable:

$$\|\psi(0) - Q_E^0\|_{H^1} \text{ small} \implies \inf_r \|\psi(t) - Q_E^0 e^{ir}\|_{H^1} \text{ small for } t \geq 0$$

(see [16]). One may expect that Q_E^0 is even *asymptotically stable* in, say, a local L^2 -norm:

$$\lim_{t \rightarrow \infty} \inf_{E, r} \|\psi(t) - Q_E^0 e^{ir}\|_{L_{loc}^2} = 0 \tag{1.4}$$

where one now has to consider the entire ground state family; i.e., to allow for modulation of the eigenvalue E as well as the phase r . If H_0 has only one discrete eigenvalue, it is proved in [18] that solutions initially close to a ground state eventually converge to some (other) ground state $e^{ir}Q_E^0$. In the case H_0 has two discrete eigenvalues $e_0 < e_1 < 0$ (also satisfying a resonance condition) and both of them are simple, this is proved in [21]. Recently, if e_1 is degenerate, the same result is proved in [9], by introducing a new type of normal form.

If the initial data is not close to a ground state, the presence of excited states makes the problem more subtle, – see [14, 20, 22]. The physical intuition is that excited states are unstable, and that, generically, nearby states should radiate, and then relax (locally) to the ground state. However, the results of [23] show that there are, in fact, *stable directions* – finite co-dimension families of solutions which converge to the excited states – at least when H_0 has simple eigenvalues. Motivated by the papers [9] and [23], in this paper we shall construct solutions converging to the nonlinear excited states of Theorem 1.1. Of course, this result does not contradict the physical intuition, since this family of solutions has zero measure in some sense (and so should not be directly seen in experiments or numerical simulations).

To study the stable directions for the nonlinear excited states, it is essential that we understand the linearized operators around these excited states, their spectral properties, and the associated time-decay estimates. In particular, if we denote the *linearized operator* around a solitary wave solution $e^{iEt}Q(x)$ by $\mathcal{L}_Q$, so that

$$\mathcal{L}_Q\zeta = -i\{(H_0 - E + 2\lambda|Q|^2)\zeta + \lambda Q^2\bar{\zeta}\},$$

then the analysis of Section 4 shows that for each of the nonlinear excited states, there is a finite-codimension subspace

$$\mathbf{E}_c(\mathcal{L}_Q) = \text{the continuous spectral subspace for } \mathcal{L}_Q$$

on which hold the dispersive decay estimates

$$\|e^{t\mathcal{L}_Q}\eta\|_{L^p} \lesssim |t|^{-3(1/2-1/p)} \|\eta\|_{L^{p'}}, \quad \frac{1}{p} + \frac{1}{p'} = 1.$$

It turns out that the spectrum of the linearized operator around Q_E (and its symmetry translates) is different from that of the linearized operator around $\tilde{Q}_E$ (and its symmetry translates). This interesting phenomena is due to the difference in the symmetry properties of Q_E and $\tilde{Q}_E$. Moreover, because of the degeneracy, there is an interaction between many different directions in the same modes, and it is more complicated to study these linearized operators, and to construct the stable directions, than it is for the ground state.

Before stating our main theorem, we make precise our further assumptions on the potential V .

Assumption A2: *The linear Hamiltonian $H_0 = -\Delta + V$ has only two eigenvalues $e_0 < e_1 < 0$, with $e_0 \neq 2e_1$.*

Remark 1.3. *Taken together, Assumptions A1 and A2 say that: the radial eigenfunction problem supports only the eigenvalue e_0 ; the eigenvalue problem corresponding to the first non-zero angular momentum sector supports only the eigenvalue e_1 ; and there are no eigenvalues corresponding to higher angular momenta. It is not difficult to construct examples of such potentials (eg. among finite square well potentials of varying depth and width).*

Next we make precise our assumptions of spatial decay and regularity of the potential $V(x)$.

Assumption A3: *For some $\sigma > 0$,*

$$|\nabla^\alpha V(x)| \leq C(1 + |x|^2)^{-5-\sigma}, \quad \forall x \in \mathbb{R}^3, \quad |\alpha| \leq 2,$$

and there is $0 < \sigma_0 < 1$ such that

$$\left\| [(x \cdot \nabla)^k V] \phi \right\|_{L^2} \leq \sigma_0 \|\Delta \phi\|_{L^2} + C \|\phi\|_{L^2}, \quad \forall k = 1, 2, 3, \quad \phi \in H^1.$$

Furthermore, 0 (the bottom of the continuous spectrum of H_0) is not an eigenvalue nor a resonance.

Assumption A3 ensures we can apply standard analysis tools for linear Schrödinger operators. It is certainly not optimal.

The final assumption will ensure, in the *resonant case* $e_0 < 2e_1$, that the resonant interaction is generic. Denote a normalized ground-state eigenfunction by ϕ_0 , which we may suppose is positive and radial (since V is radial), denote by $\mathbf{P}_0(H_0)$ the corresponding orthogonal projection,

$$\text{Null}(H_0 - e_0) = \mathbb{C}\phi_0, \quad \phi_0(x) = \phi_0(|x|) > 0, \quad \mathbf{P}_0 = \mathbf{P}_0(H_0) = \langle \phi_0 | \cdot \rangle \langle \cdot | \phi_0 \rangle,$$

and denote by $\mathbf{P}_c$ the orthogonal projection onto the continuous spectral subspace of H_0 :

$$\mathbf{P}_c = \mathbf{P}_c(H_0) = 1 - \mathbf{P}_0(H_0) - \mathbf{P}_1(H_0).$$

Assumption A4: *“Fermi Golden Rule”: If $e_0 < 2e_1$ (“resonant case”), there exists $\lambda_0 > 0$ such that*

$$\lim_{r \rightarrow 0^+} \text{Im} \left(\phi_0 \phi^2, (H_0 - 2e_1 + e_0 - ir)^{-1} \mathbf{P}_c \phi_0 \phi^2 \right) \geq \lambda_0 \|\phi\|_{L^2}^4 \quad \forall \phi \in \mathbf{V}. \quad (1.5)$$

When e_1 is simple, the condition (1.5) is well-known and appears in many papers, see [3, 4, 5, 8, 14, 20, 21, 22, 23, 24]. In the degenerate case, (1.5) is also used in [9], and it is claimed there that (1.5) holds generically.

The existence of stable directions for the excited states is proved in Section 5:

Theorem 1.2 (Stable directions for degenerate excited states). *Assume that $H_0 = -\Delta + V$ satisfies A1-A4 above. Then, there exists $\epsilon_0 > 0$ such that for any $0 < \epsilon \leq \epsilon_0$, there is $\delta_0 > 0$ such that for any $0 < \delta < \delta_0$ the following holds: if Q denotes a nonlinear excited state with $E - e_1 = \lambda\epsilon^2$, $\mathcal{L}_Q$ denotes the corresponding linearized operator, and $\eta_\infty \in W^{2,1} \cap H^2 \cap \mathbf{E}_c(\mathcal{L}_Q)$ with $\|\eta_\infty\|_{W^{2,1} \cap H^2} \leq \delta$, then there exists a solution $\psi(x, t)$ of (1.1) such that*

$$\|\psi(x, t) - \psi_{as}(x, t)\|_{H^2} \leq C(\epsilon)\delta^{7/4}(1+t)^{-1},$$

where

$$\psi_{as}(x, t) := e^{-iEt}[Q + e^{t\mathcal{L}_Q}\eta_\infty]$$

and so in particular, for $p > 2$,

$$\|\psi(x, t) - e^{-iEt}Q(x)\|_{L^p} \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

2 Some notation and definitions

- (1) Let $L^2 := L^2(\mathbb{R}^3, \mathbb{C})$, $\mathbf{E} := L^2(\mathbb{R}^3, \mathbb{C}^2)$. We equip $\mathbf{E}$ with the standard inner product

$$\langle f, g \rangle = \int_{\mathbb{R}^3} (\bar{f}_1 g_1 + \bar{f}_2 g_2) dx, \quad \forall f = \begin{bmatrix} f_1 \\ f_2 \end{bmatrix}, \quad g = \begin{bmatrix} g_1 \\ g_2 \end{bmatrix} \in \mathbf{E}.$$

Moreover, for any $u \in L^2$, we shall write $\vec{u} = \begin{bmatrix} \text{Re}(u) \\ \text{Im}(u) \end{bmatrix} \in \mathbf{E}$.

- (2) Recall

$$\begin{aligned} H_0 &= -\Delta + V \\ \mathbf{V} &:= \text{Null}(H_0 - e_1) = \text{span}_{\mathbb{C}}\{\phi_1, \phi_1, \phi_2\}, \quad \text{Null}(H_0 - e_0) = \mathbb{C}\phi_0 \\ \mathbf{P}_0 &= \langle \phi_0 \mid, \quad \mathbf{P}_1 = \sum_{j=1}^3 \langle \phi_j \mid, \quad \mathbf{P}_c = 1 - P_0 - P_1 \end{aligned}$$

(3) Denote

$$J = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad \sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (2.1)$$

(4) Let $\mathbf{O}(k)$ denote the group of orthogonal transformations on $\mathbb{R}^k$. Identifying $\mathbb{R}^2$ with $\mathbb{C}$, we can write $\mathbf{O}(2)$ as the group generated by $\{e^{ir}, \mathbf{conj} : r \in [0, 2\pi)\}$, where $\mathbf{conj}$ denotes complex conjugation.

Definition 2.1. Let $\mathbf{G} := \mathbf{O}(3) \oplus \mathbf{O}(2)$. For $g = (g_1, g_2) \in G$, define its action on L^2 by $g * f(x) := g_2 * f(g_1 * x)$ where $g_1 * x$ denotes usual matrix multiplication, and $g_2 *$ denotes complex multiplication (or conjugation).

(5) For $\alpha \in [0, 2\pi)$, we denote by $R_{jk}(\alpha)$ the rotation matrix in the $x_j x_k$ -plane of $\mathbb{R}^3$ through angle α , for $j, k = 1, 2, 3$, and $j < k$. Precisely,

$$\begin{aligned} R_{12}(\alpha) &:= \begin{bmatrix} \cos(\alpha) & -\sin(\alpha) & 0 \\ \sin(\alpha) & \cos(\alpha) & 0 \\ 0 & 0 & 1 \end{bmatrix}, & R_{13}(\alpha) &:= \begin{bmatrix} \cos(\alpha) & 0 & \sin(\alpha) \\ 0 & 1 & 0 \\ -\sin(\alpha) & 0 & \cos(\alpha) \end{bmatrix}, \\ R_{23}(\alpha) &:= \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\alpha) & -\sin(\alpha) \\ 0 & \sin(\alpha) & \cos(\alpha) \end{bmatrix}. \end{aligned} \quad (2.2)$$

Also, let

$$\begin{aligned} \Gamma(\delta, \alpha, \sigma) &:= R_{12}(\delta)R_{13}(\alpha)R_{23}(\sigma), & \Gamma_0(\alpha, \delta) &:= R_{12}(\alpha)R_{13}(\delta), \\ \Gamma_1(\alpha, \delta) &:= R_{13}(\alpha)R_{23}(\delta), & \alpha, \delta, \sigma &\in [0, 2\pi). \end{aligned} \quad (2.3)$$

By the Euler-Brauer resolution of a rotation, for any rotation matrix $A \in \mathbf{SO}(3)$, there exist unique $(\delta, \alpha, \sigma) \in [0, 2\pi)^3$ such that $A = \Gamma(\delta, \theta, \sigma)$ (eg. [15, p. 146]). Moreover, for any $B \in \mathbf{O}(3)$, either B or $-B$ lies in $\mathbf{SO}(3)$.

(6) For some $s > 3$, let L_s^2 be the weighted L^2 space defined by

$$L_s^2 := \{f : (1 + |x|^2)^{s/2} f(x) \in L^2(\mathbb{R}^3, \mathbb{C})\}. \quad (2.4)$$

Then, let $\mathbf{E}_s = L_s^2 \times L_s^2$ and $\mathbf{B} := \mathbf{B}(\mathbf{E}_s, \mathbf{E}_{-s})$ be the space of all bounded operators from $\mathbf{E}_s$ to $\mathbf{E}_{-s}$.

3 Existence of Nonlinear Excited States

Let $\mu := E - e_1$ and

$$F(\mu, Q) := (H_0 - e_1)Q + \lambda|Q|^2Q - \mu Q. \quad (3.1)$$

Since $V = V(|x|)$, we see that F is invariant under the action the group $\mathbf{G}$, i.e.

$$F(\mu, g * Q) = g * F(\mu, Q), \quad \forall g \in \mathbf{G}.$$

As remarked in the introduction, the equation $F(\mu, Q) = 0$ has a solution for Q near $\mathbf{V}$ only if $\lambda\mu = \lambda(E - e_1) > 0$. Now, we write $Q = \epsilon(v + h)$ where we will take $\epsilon = \sqrt{\lambda\mu} > 0$ sufficiently small, $v \in \mathbf{V}$ of order one, and $h \in \mathbf{V}^\perp$. Then the equation $F(\mu, Q) = 0$ becomes

$$(H_0 - e_1)h + \lambda\epsilon^2|v + h|^2 - \mu(v + h) = 0. \quad (3.2)$$

Now, applying the projections $\mathbf{P}_1$ and $\mathbf{P}_1^\perp := 1 - \mathbf{P}_1$ to the equation (3.2), we get

$$h = \lambda\epsilon^2[H_0 - e_1]^{-1}\mathbf{P}_1^\perp\{h - |v + h|^2(v + h)\} \quad (3.3)$$

$$\mathbf{P}_1[v - |v + h|^2(v + h)] = 0. \quad (3.4)$$

By applying the contraction mapping theorem or implicit function theorem, we see that for any fixed $c_1 > 0$, there exists a sufficiently small number $\epsilon_1 > 0$, such that for all $0 \leq \epsilon < \epsilon_1$, and each $v \in \mathbf{V}$ with $\|v\| < c_1$, there is a unique solution $h = h(\epsilon, v) \in H^2$ of (3.3) satisfying

$$h(0, v) = 0, \quad h_v(0, v) = 0.$$

Moreover, since $\mathbf{P}_1 g = g \mathbf{P}_1$ for all $g \in \mathbf{G}$, by the uniqueness of the solution h of (3.3), we also have

$$h(\epsilon, g * v) = g * h(\epsilon, v), \quad \forall g \in \mathbf{G}. \quad (3.5)$$

Let

$$N(\epsilon, v) := \mathbf{P}_1[v - |v + h(\epsilon, v)|^2(v + h(\epsilon, v))].$$

Then, we have

$$N(\epsilon, g * v) = g * N(\epsilon, v), \quad \text{for all } v \in \mathbf{V} \text{ and } g \in \mathbf{G}. \quad (3.6)$$

Moreover,

$$N_v(0, v)w = \mathbf{P}_1[w - 2|v|^2w - v^2\bar{w}], \quad \forall w \in \mathbf{V}. \quad (3.7)$$

In order to apply the implicit function theorem to solve the equation $N(\epsilon, v) = 0$, we need to find $v_0 \in \mathbf{V}$ such that

$$N(0, v_0) = 0, \quad \text{and} \quad N_v(0, v_0) : \mathbf{V} \rightarrow \mathbf{V} \text{ is invertible.} \quad (3.8)$$

Set $v_0 = \phi_z := z \cdot \phi = z_1\phi_1 + z_2\phi_2 + z_3\phi_3$ for $z \in \mathbb{C}^3$. Then, $N(0, v_0) = 0$ if and only if $(\phi_j, N(0, \phi_z)) = 0$ for all $j = 1, 2, 3$. Set $I = (\phi_1^2, \phi_2^2)$. For each $j = 1, 2, 3$, the equation $(\phi_j, N(0, \phi_z)) = 0$ becomes

$$\begin{aligned} z_j &= (\phi_j, |\phi_z|^2 \phi_z) = 2z_j \sum_{l \neq j} |z_l|^2 (\phi_j^2, \phi_l^2) + \bar{z}_j \sum_l z_l^2 (\phi_j^2, \phi_l^2) \\ &= (\phi_1^2, \phi_2^2) \left\{ 3|z_j|^2 z_j + \bar{z}_j \sum_{l \neq j} z_l^2 + 2z_l \sum_{l \neq j} |z_l|^2 \right\} \\ &= I \{ 2z_j |z|^2 + \bar{z}_j z^2 \}, \quad \text{where} \quad z^2 := \sum_l z_l^2. \end{aligned}$$

Equivalently,

$$2z_j |z|^2 + \bar{z}_j z^2 = \frac{1}{I} z_j \quad \forall j = 1, 2, 3. \quad (3.9)$$

Write $z_j = a_j e^{i\alpha_j}$, for $a_j \geq 0$ and $\alpha_j \in [0, 2\pi)$. For some j_0 , we have $z_{j_0} \neq 0$, and by applying a phase rotation we may assume $z_{j_0} \in \mathbb{R}$. Then dividing (3.9) by z_{j_0} , we get

$$2|z|^2 + z^2 = \frac{1}{I}. \quad (3.10)$$

Moreover for any j with $z_j \neq 0$, (3.9) implies

$$2|z|^2 + e^{-2i\alpha_j} z^2 = \frac{1}{I}, \quad (3.11)$$

and so, comparing (3.10) and (3.11), we see that either $z_j = \pm a_j$, or else $z^2 = 0$. So (3.9) is equivalent to

$$|z|^2 = \frac{1}{2I} \quad \text{and} \quad z^2 = 0; \quad \text{or} \quad |z|^2 = \frac{1}{3I} \quad \text{and} \quad e^{i\alpha} z \in \mathbb{R}^3, \text{ some } \alpha \in [0, 2\pi). \quad (3.12)$$

Thus we obtain two representative elements of the solutions of $N(0, v) = 0$:

Lemma 3.1. *Let $v_1 = (1/3I)^{1/2}(1, 0, 0) \cdot \phi$, $v_2 = (1/4I)^{1/2}(1, i, 0) \cdot \phi$, and let $\mathbf{O}_1, \mathbf{O}_2$ be respectively the orbits of v_1 and v_2 under the action of $\mathbf{G}$ on $\mathbf{V}$. The set $\mathbf{O} := \mathbf{O}_1 \cup \mathbf{O}_2$ contains all non-zero solutions of $N(0, v) = 0$.*

Proof. First of all, note that $\mathbf{V}$ is invariant under the action of $\mathbf{G}$. So, the action of $\mathbf{G}$ on $\mathbf{V}$ is well-defined. Moreover, from (3.6), we see that v solves the equation $N(0, v) = 0$ for all $v \in \mathbf{O}$. Now observe that for any $v = z \cdot \phi \in \mathbf{V}$, we have

$$g * v = g_2 * [z' \cdot \phi], \quad \forall g = (g_1, 1) \in \mathbf{G}, \quad z' := g_1 z. \quad (3.13)$$

So, to prove that for every $v = z \cdot \phi$ with $z = (z_1, z_2, z_3) \in \mathbf{C}^3$ such that $N(0, v) = 0$ then $v \in \mathbf{O}$, we shall need to find some $g = (g_1, g_2) \in \mathbf{G}$ such that

$$g_2 * [z' \cdot \phi] = v_1 \text{ or } = v_2, \quad \text{with } z' := g_1 z.$$

So let $v = z \cdot \phi$ with $z = (z_1, z_2, z_3) \in \mathbf{C}^3$ such that $N(0, v) = 0$. Then, z satisfies (3.12). If $e^{i\alpha} z \in \mathbb{R}^3$ for some $\alpha \in [0, 2\pi)$ with $|z|^2 = (1/3I)^{1/2}$, then, it's simple to see that there exists $g := (g_1, e^{-i\alpha}) \in \mathbf{G}$ with $g_1 \in \mathbf{SO}(3)$ such that $g * v_1 = z \cdot \phi$. So, $z \in \mathbf{O}_1$. Now, assume that $e^{i\alpha} z \notin \mathbb{R}^3$ for all $\alpha \in [0, 2\pi)$. Then, we have $|z|^2 = 1/(2I)$ and $z^2 = 0$. We write $z_j = a_j e^{i\alpha_j}$, for $a_j \geq 0$ and $0 \leq \alpha_j < 2\pi$ for all $j = 1, 2, 3$. Applying a spatial rotation, and then a phase rotation, we may assume that $a_1 > 0$ and $\alpha_1 = 0$. So, $z = (a_1, a_2 e^{i\alpha_2}, a_3 e^{i\alpha_3})$. Now, if $a_2 a_3 = 0$ or $\alpha_2 \alpha_3 = 0$, then it is also simple to show that $v \in \mathbf{O}$. So, we assume that $a_1 a_2 a_3 \neq 0$ and $\alpha_2 \alpha_3 \neq 0$. Let $M_1 := R_{23}(\alpha)^T$ where R_{23} is defined in (2.2) and choose α so that $a_2 \cos(\alpha) \cos(\alpha_2) = a_2 \sin(\alpha) \cos(\alpha_3)$, or in other words $\tan(\alpha) = a_2 \cos(\alpha_2) / [a_3 \cos(\alpha_3)]$. Then, $(M_1, 1) \in \mathbf{G}$ and $(M_1, 1) * v = v' := z' \cdot \phi$ where $z' := (r_1, ia'_2, a'_3 e^{i\alpha'_3})$ for some $a'_2 \in \mathbb{R}$ and $a'_3 \geq 0$. Since $z^2 = 0$ and $|z|^2 = 1/(2I)$, we have $(z')^2 = 0$ and $|z'| = 1/(2I)$. So,

$$a_1 - (a'_2)^2 + (a'_3)^2 e^{2i\alpha'_3} = 0.$$

This implies that $e^{2i\alpha'_3} \in \mathbb{R}$. Therefore, $\alpha'_3 \in \{0, \pi/2, \pi, 3\pi/2\}$. So either $z' = (r_1, ia, ib)$ or $z' = (r_1, ia, b)$ for some $a, b \in \mathbb{R}$ such that $|z'| = 1/(2I)$. From this and by taking $M_2 = R_{23}(\theta')^T$ or $M_2 = R_{13}(\theta')^T$ with appropriate θ' , we see that $(M_2, 1)v' = v'' := z'' \cdot \phi$ with $z'' = (c, id, 0)$ for some $c, d \in \mathbb{R}, c \neq 0$ and $|z''|^2 = 1/(2I)$ and $(z'')^2 = 0$. Then, it follows that $d \neq 0$ and then $v'' \in \mathbf{O}_2 \subset \mathbf{O}$. This completes the proof of the lemma. $\square$

Lemma 3.1 completely solves the first equation of (3.8). For the second condition in (3.8), it is noted that

$$N(\epsilon, v)(iv) = 0, \quad \forall v \in \mathbf{V} : N(\epsilon, v) = 0,$$

a consequence of the phase invariance. So, the second condition of (3.8) never holds. To overcome this and solve the equation $N(\epsilon, v) = 0$, we shall restrict N to the invariant subspaces of $\mathbf{V}$ and solve $N = 0$ on these subspaces. To this end, we introduce the following lemma:

Lemma 3.2. *For each $j = 1, 2$, let $\mathbf{V}_j := \text{span}_{\mathbb{R}}\{v_j\}$ where v_1, v_2 are defined in Lemma 3.1. There are subgroups $\mathbf{G}_j \leq \mathbf{G}$ such that $\mathbf{V}_j$ is the fixed subspace of $\mathbf{V}$ under the action of $\mathbf{G}_j$ on $\mathbf{V}$. In other words,*

$$\mathbf{V}_j = \{v \in \mathbf{V} : g * v = v \quad \forall g \in \mathbf{G}_j\}, \quad j = 1, 2. \quad (3.14)$$

Proof. Recall that $\mathbf{conj}$ denotes complex conjugation: $\mathbf{conj} * z = \bar{z}$ for $z \in \mathbb{C}$. Set

$$\iota_0 := \left(\begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, -1 \right), \quad \iota_1 := \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \mathbf{conj} \right), \quad \iota_2 := \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, 1 \right) \in \mathbf{G}.$$

Moreover, let

$$g(\alpha) := (R_{12}(\alpha), e^{-i\alpha}) \in \mathbf{G} \quad \text{and} \quad G'_1 := \left\{ \begin{bmatrix} 1 & 0 \\ 0 & g \end{bmatrix}, g \in \mathbf{O}(2) \right\} \leq \mathbf{O}(3).$$

Then, let $\mathbf{G}_1$ be the subgroup of $\mathbf{G}$ which is generated by the subgroup $\mathbf{G}'_1 \oplus \{1, \mathbf{conj}\}$ and $\{\iota_0\}$. And let $\mathbf{G}_2$ be the subgroup of $\mathbf{G}$ which is generated by $\{\iota_1, \iota_2, g(\alpha), \forall \alpha \in [0, 2\pi)\}$. We shall show that $\mathbf{V}_j$ is the fixed subspace of the action of $\mathbf{G}_j$ on $\mathbf{V}$ for $j = 1, 2$. Note that

$$\{v \in \mathbf{V} : (g, 1) * v = v \text{ and } \iota_0 * v = v, \forall g \in \mathbf{G}'_1\} = \text{span}_{\mathbb{C}}\{\phi_1\}.$$

Moreover, for all $z \in \mathbb{C}$, we see that $\mathbf{conj} * z\phi_1 = z\phi_1$ if and only if $z \in \mathbb{R}$. So, we obtain (3.14) for $j = 1$. On the other hand, for $v = z \cdot \phi \in \mathbf{V}$ such that v is fixed by the action of $\mathbf{G}_2$, we have $g(\alpha) * v = v$ for all $\alpha \in [0, 2\pi)$. So,

$$\begin{cases} e^{i\alpha} z_1 &= z_1 \cos(\alpha) + z_2 \sin(\alpha), \\ e^{i\alpha} z_2 &= -z_1 \sin(\alpha) + z_2 \cos(\alpha), \\ e^{i\alpha} z_3 &= z_3, \quad \forall \alpha \in [0, 2\pi). \end{cases}$$

Therefore, we get $z_2 = iz_1$ and $z_3 = 0$. So $v = z_1(\phi_1 + i\phi_2)$. Moreover, $\iota_1 * v = v, \iota_2 * v = v$ imply that $z_1 \in \mathbb{R}$. So, $v \in \mathbf{V}_2$ and this completes the proof of the lemma. $\square$

An elementary but important observation: (3.6) implies that $N = N(\epsilon, \cdot)$ maps $\mathbf{V}_j$ into $\mathbf{V}_j$. Thus we have well-defined maps

$$N_1 := N|_{\mathbf{V}_1} : \mathbf{V}_1 \rightarrow \mathbf{V}_1, \quad N_2 := N|_{\mathbf{V}_2} : \mathbf{V}_2 \rightarrow \mathbf{V}_2.$$

Now consider the bifurcation equation $N_1 = 0$. From the definition of v_1 in Lemma 3.1, we have $N_1(0, v_1) = 0$. Moreover, from (3.7)

$$\partial_v N_1(0, v_1)(\phi_1) = (\phi_1, N_v(0, v_1)\phi_1)\phi_1 = -2\phi_1.$$

Therefore, applying the implicit function theorem, we obtain a bifurcation solution of the equation $N_1 = 0$ as

$$v_+(\epsilon) = v_1 + a_+(\epsilon)\phi_1, \quad a_+(\epsilon) = O(\epsilon) \quad (3.15)$$

for $0 < \epsilon < \epsilon_2$, some $0 < \epsilon_2 \leq \epsilon_1$. Now, we consider the bifurcation equation $N_2 = 0$. Let $\phi^* = \phi_1 + i\phi_2$. As before, we have $N_2(0, v_2) = 0$ and

$$\partial_v N_2(0, v_2)(\phi^*) = (\phi^*, N_v(0, v_2)\phi^*)\phi^* = [2 - \frac{3}{4I}(|\phi^*|^2, |\phi^*|^2)]\phi^* = -4\phi^*.$$

Again, by the implicit function theorem, we have a bifurcation solution of the equation $N_2 = 0$ given by

$$v_-(\epsilon) = v_2 + a_-(\epsilon)\phi^*, \quad a_-(\epsilon) = O(\epsilon) \quad (3.16)$$

for $0 < \epsilon < \epsilon_2$, some $0 < \epsilon_2 \leq \epsilon_1$.

We denote the two resulting solutions of (3.1) by

$$\begin{aligned} Q_E &:= \epsilon[v_1 + a_+(\epsilon)\phi_1 + h(\epsilon, v_1)] = \rho(\epsilon)\phi_1 + h(\epsilon, v_1), \\ \tilde{Q}_E &:= \epsilon[v_2 + a_-(\epsilon)\phi^* + h(\epsilon, v_2)] = \tilde{\rho}(\epsilon)\phi^* + h(\epsilon, v_2), \end{aligned} \quad (3.17)$$

where $\rho(\epsilon) := \epsilon/(3I)^{1/2} + \epsilon a_+(\epsilon)$ and $\tilde{\rho}(\epsilon) := \epsilon/(4I)^{1/2} + \epsilon a_-(\epsilon)$.

Remark 3.1. By (3.5), we see that for each $j = 1, 2$, we have $h(\epsilon, v_j) = h(\epsilon, g * v_j) = g * h(\epsilon, v_j)$ for all $g \in \mathbf{G}_j$. Therefore, we can write Q_E and $\tilde{Q}_E$ as

$$Q_E = x_1 f_1(x_1^2, x_2^2 + x_3^2), \quad \tilde{Q}_E = e^{i\theta} f_2(x_1^2 + x_2^2, x_3^2)$$

for some real functions f_1, f_2 . Here, $\theta \in [0, 2\pi)$ is the angle between that x_1 axis and the vector $x' = (x_1, x_2) \in \mathbb{R}^2$.

Note that $Q_E, \tilde{Q}_E \in H^2$ by construction. Their exponential decay is standard; in particular, the argument in the case of simple eigenvalues – see, eg, [12] – applies. This completes the proof of Theorem 1.1. $\square$

Remark 3.2. It is straightforward to express the full solution branches explicitly in terms of symmetry transformations:

$$Q(x) = e^{ir_1} Q_E(\Gamma_0(r_2, r_3)x) \quad \text{and} \quad \tilde{Q}(x) = e^{ir_1} \tilde{Q}_E(\Gamma_1(r_2, r_3)x) \quad (3.18)$$

for parameters $E \in (e_1, e_1 + \lambda\epsilon_2^2)$, $r_1, r_2, r_3 \in [0, 2\pi)$.

4 Spectral Analysis of the Linearized Operators Around Excited States

The spectral properties are invariant under symmetry transformations of the underlying solution, hence we may fix one element of each branch of excited states. So in what follows, let $Q = Q_E$ or $\tilde{Q}_E$. We write solution $\psi(t, x)$ of (1.1) as $\psi(t, x) = [Q + \zeta]e^{-iEt}$. Then, from (1.1), we have

$$\partial_t \zeta = \mathcal{L}_Q \zeta + \text{nonlinear terms},$$

where

$$\mathcal{L}_Q \zeta = -i\{(H_0 - E + 2\lambda|Q|^2)\zeta + \lambda Q^2 \bar{\zeta}\}. \quad (4.1)$$

Decomposing ζ into the real and imaginary parts, we can extend $\mathcal{L}$ to $\vec{\mathcal{L}} : \mathbf{E} \rightarrow \mathbf{E}$. That is,

$$\vec{\mathcal{L}}_Q := J(H_0 - E) + W_Q, \quad \text{where} \quad J := \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad W_Q := \begin{bmatrix} W_{1,Q} & W_{2,Q} \\ W_{3,Q} & W_{4,Q} \end{bmatrix}, \quad (4.2)$$

with

$$\begin{aligned} W_{1,Q} &:= 2\lambda \operatorname{Re} Q \operatorname{Im} Q, & W_{4,Q} &= -W_{1,Q}, \\ W_{2,Q} &:= \lambda(2|Q|^2 + [\operatorname{Im} Q]^2 - [\operatorname{Re} Q]^2), \\ W_{3,Q} &:= -\lambda(2|Q|^2 + [\operatorname{Re} Q]^2 - [\operatorname{Im} Q]^2). \end{aligned}$$

For $Q = Q_E$ we will write $\mathbf{L}_0 := \vec{\mathcal{L}}_{Q_E}$ and for $Q = \tilde{Q}_E$, we will write $\mathbf{H}_0 := \vec{\mathcal{L}}_{\tilde{Q}_E}$. So

$$\begin{aligned} \mathbf{L}_0 &= \begin{bmatrix} 0 & L_- \\ -L_+ & 0 \end{bmatrix}, \quad \text{with} \quad L_{\pm} := H_0 - E + \lambda(2 \pm 1)Q_E^2, \\ \mathbf{H}_0 &= J(H_0 - E) + \begin{bmatrix} W_1 & W_2 \\ W_3 & W_4 \end{bmatrix}, \quad W_j := W_{j, \tilde{Q}_E}. \end{aligned} \quad (4.3)$$

We now state several lemmas which will be used in the next sections. These lemmas closely parallel lemmas in [23], and so we will omit some details.

Lemma 4.1. *Let $\Sigma_c := \{ia : |a| \geq |E|\}$. For all $\tau \in \Sigma_c$, $|\tau| \neq |E|$, the equation $\vec{\mathcal{L}}\psi = \tau\psi$ has no non-zero solution $\psi \in \mathbf{E}$. Moreover, $\pm i|E|$ are neither eigenvalues nor resonances.*

Proof. Note that $\vec{\mathcal{L}} = J(-\Delta - E) + \widetilde{W}$ where $\widetilde{W} = JV + W$. By the assumptions on V , and the exponential decay of Q , the potential $\widetilde{W}$ decays quickly. The first part of the lemma then follows as in [23, Section 2.4], making use of the resolvent estimate Lemma 4.17. To prove that $\pm i|E|$ is not an eigenvalue or resonance, we may use Lemma 4.17 and the argument in [13], or we can follow [23, Section 2.5]. $\square$

Lemma 4.2. *Let $R(z) := (\vec{\mathcal{L}} - z)^{-1}$, $R_0(z) = [JH - z]^{-1}$ and $\Sigma_p := \{0, \pm i(e_0 - E)\}$ where $H = H_0 - E$. There exists an order one constant $C > 0$ such that*

$$\|R(z)\|_{(\mathbf{E}, \mathbf{E})} \leq C[\|R_0(z)\|_{(\mathbf{E}, \mathbf{E})} + \|R_0(z)\|_{(\mathbf{E}, \mathbf{E})}^2], \quad \forall z \notin \Sigma_c : \text{dist}(z, \Sigma_p) \geq \epsilon.$$

Proof. We have

$$R(z) = [1 + R_0(z)W]^{-1}R_0(z) = \sum_{j=0}^{\infty} (-1)^j [R_0(z)W]^j R_0(z).$$

Using this and the facts that $R_0(z)$ is uniformly bounded in $\mathbf{B}$ with bound $O(1/\epsilon)$ for z with $\text{dist}(z, \Sigma_p) \geq \epsilon$ (see [13]), and W is localized of order ϵ^2 , we obtain

$$\begin{aligned} \|R(z)\|_{(\mathbf{E}, \mathbf{E})} &\leq \|R_0(z)\|_{(\mathbf{E}, \mathbf{E})} + \sum_{j=1}^{\infty} C \|R_0(z)\|_{(\mathbf{E}, \mathbf{E})} \{C\epsilon^2 \|R_0(z)\|_{\mathbf{B}}\}^{j-1} \|R_0(z)\|_{(\mathbf{E}, \mathbf{E})} \\ &\leq C \left\{ \|R_0(z)\|_{(\mathbf{E}, \mathbf{E})} + \|R_0(z)\|_{(\mathbf{E}, \mathbf{E})}^2 \sum_{j=0}^{\infty} [C\epsilon]^j \right\} \end{aligned}$$

For ϵ sufficiently small, the series in the right hand side of the last inequality converges, and the lemma follows. $\square$

Lemma 4.3. *Let $\mathbf{D}_1 := \{a + ib : |b - (e_0 - E)| \leq \epsilon, 0 < a \leq \epsilon\}$ and $R_0(z) = (JH - z)^{-1}$. Then for some fixed $s > 3$, there is a constant $C > 0$ such that for all $z \in \mathbf{D}_1$,*

$$\begin{aligned} \|\langle x \rangle^{-s} \mathbf{P}_c R_0(z) \mathbf{P}_c \langle x \rangle^{-s}\|_{(L^2, L^2)} &\leq C, \\ \left\| \langle x \rangle^{-s} \mathbf{P}_c \frac{d}{dz} R_0(z) \mathbf{P}_c \langle x \rangle^{-s} \right\|_{(L^2, L^2)} &\leq C(\text{Re } z)^{-1/2}. \end{aligned} \tag{4.4}$$

Here $\mathbf{P}_c = \mathbf{P}_c(JH) = \begin{bmatrix} \mathbf{P}_c(H) \\ \mathbf{P}_c(H) \end{bmatrix}$. Moreover, for $z_1, z_2 \in \mathbf{D}_1$, we have

$$\|\langle x \rangle^{-s} \mathbf{P}_c [R_0(z_1) - R_0(z_2)] \mathbf{P}_c \langle x \rangle^{-s}\|_{(L^2, L^2)} \leq C[\max\{\text{Re } z_1, \text{Re } z_2\}]^{-1/2} |z_1 - z_2|. \tag{4.5}$$

Proof. We write $R_0(z) = R_{01}(z) + R_{02}(z)$, where

$$R_{01}(z) := \frac{1}{2} \begin{bmatrix} i & -1 \\ 1 & i \end{bmatrix} (H - iz)^{-1}, \quad R_{02}(z) := \frac{1}{2} \begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix} (H + iz)^{-1}. \quad (4.6)$$

From this and since R_{02} is regular in $\mathbf{D}_1$, we shall prove the lemma with $R_0(z)$ replaced by $T_0(z) := (H - iz)^{-1} \mathbf{P}_c(H)$. The first estimate in (4.4) is standard and therefore we skip its proof. The second estimate in (4.4) follows from (4.5). So, it's sufficient to prove (4.5). The proof now is similar to that of [23, Lemma 2.3] and we shall only give the main steps.

For any z_1, z_2 in $\mathbf{D}_1$. We write $z_1 = a_1 + ib_1, z_2 = a_2 + ib_2$ and assume that $a_1 \leq a_2$. Let $z_3 = a_2 + ib_1$. We have $z_3 \in \mathbf{D}_1$. For any $u, v \in L^2$ with $\|u\|_2 = \|v\|_2 = 1$, let $u_1 = \mathbf{P}_c \langle x \rangle^{-s} u$, $v_1 = \mathbf{P}_c \langle x \rangle^{-s} v$. We have $u_1, v_1 \in L^1 \cap L^2$. Now, using the decay estimate $\|e^{-itH} \mathbf{P}_c \phi\|_{L^\infty} \leq C(1+t)^{-3/2} \|\phi\|_{L^1}$ with $H = H_0 - E$, we get

$$\begin{aligned} & |(u, \langle x \rangle^{-s} \mathbf{P}_c [T_0(z_1) - T_0(z_3)] \mathbf{P}_c \langle x \rangle^{-s} v)| \\ &= \left| \int_0^\infty (u_1, [e^{-it(H-i(a_1+ib_1))} - e^{-it(H-i(a_2+ib_1))}] v_1) dt \right| \\ &= \left| \int_0^\infty (u_1, e^{-it(H+ib_1)} v_1) (e^{-a_1 t} - e^{-a_2 t}) dt \right| \\ &\leq C \int_0^\infty (1+t)^{-3/2} (e^{-a_1 t} - e^{-a_2 t}) dt \leq C a_2^{-1/2} (a_2 - a_1). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} & |(u, \langle x \rangle^{-s} \mathbf{P}_c [T_0(z_3) - T_0(z_2)] \mathbf{P}_c \langle x \rangle^{-s} v)| \\ &= \left| \int_0^\infty (u_1, [e^{-it(H-i(a_2+ib_1))} - e^{-it(H-i(a_2+ib_2))}] v_1) dt \right| \\ &= \left| \int_0^\infty (u_1, e^{-it(H+ib_2-ia_2)} v_1) (e^{it(b_2-b_1)} - 1) dt \right| \\ &\leq C \int_0^\infty (1+t)^{-3/2} e^{-ta_2} |e^{it(b_2-b_1)} - 1| dt \leq C a_2^{-1/2} |b_2 - b_1|. \end{aligned}$$

Therefore,

$$\begin{aligned} & |(u, \langle x \rangle^{-s} \mathbf{P}_c [T_0(z_1) - T_0(z_2)] \mathbf{P}_c \langle x \rangle^{-s} v)| \\ &\leq |(u, \langle x \rangle^{-s} \mathbf{P}_c [T_0(z_1) - T_0(z_3)] \mathbf{P}_c \langle x \rangle^{-s} v)| + |(u, \langle x \rangle^{-s} \mathbf{P}_c [T_0(z_3) - T_0(z_2)] \mathbf{P}_c \langle x \rangle^{-s} v)| \\ &\leq C a_2^{-1/2} (|a_1 - a_2| + |b_1 - b_2|) \leq C \operatorname{Re}(z_2)^{-1/2} |z_1 - z_2|. \end{aligned}$$

This completes the proof of the lemma. $\square$

4.1 Spectral Analysis of $\mathbf{L}_0$

We study the spectrum of $\mathbf{L}_0$ first. From (2.1) and (4.3), we have

$$\sigma_1 \mathbf{L}_0 = \mathbf{L}_0^* \sigma_1, \quad \sigma_3 \mathbf{L}_0 = -\mathbf{L}_0 \sigma_3, \quad \text{where} \quad \mathbf{L}_0^* = \begin{bmatrix} 0 & -L_+ \\ L_- & 0 \end{bmatrix}, \quad (4.7)$$

Lemma 4.4 (Spectrum of L_-). *For $Q = Q_E$, the following hold:*

- (i) *The discrete spectrum of L_- consists of $\{\tilde{e}_0, \tilde{e}_2, 0\}$ where 0 and $\tilde{e}_0 = e_0 - e_1 + O(\epsilon^2)$ are simple eigenvalues, and $\tilde{e}_2 = \tilde{e}_3 = -2\lambda\epsilon^2/3 + O(\epsilon^3)$ is a double eigenvalue. For $j = 0, 1, 2, 3$, there exist localized orthonormal functions $\tilde{\phi}_j = \phi_j + O(\epsilon^2)$ such that*

$$L_- \tilde{\phi}_0 = \tilde{e}_0 \tilde{\phi}_0, \quad L_- \tilde{\phi}_1 = 0, \quad L_- \tilde{\phi}_j = \tilde{e}_j \tilde{\phi}_j, \quad j = 2, 3.$$

Moreover, $\tilde{\phi}_0(x)$ is even with respect to x_2, x_3 and $\tilde{\phi}_j(x)$ are odd with respect to x_j and even with respect to x_k for all $j, k > 1$ and $j \neq k$.

- (ii) *The continuous spectrum of L_- is $\sigma_c(L_-) = [|E|, \infty)$.*

Proof. We shall just prove (i), as (ii) is standard. Since L_- is a small perturbation of $H = H_0 - E$, they have the same number of discrete eigenvalues. From Assumption A0, we see that H has a simple eigenvalue $e_0 - E$ with eigenfunction ϕ_0 and a triple eigenvalue $e_1 - E$ with eigenfunctions ϕ_1, ϕ_2, ϕ_3 . We will compute the discrete eigenvalues of L_- which are perturbations of the discrete eigenvalues of H .

Since $e_0 - E$ is simple, by standard perturbation theory, we can find an eigenfunction $\tilde{\phi}_0 = \phi_0 + O(\epsilon^2)$ such that $L_- \tilde{\phi}_0 = \tilde{e}_0 \tilde{\phi}_0$ and $\tilde{e}_0 = e_0 - E + O(\epsilon^2) = e_0 - e_1 + O(\epsilon^2)$. Moreover, direct computation gives $L_- Q = 0$. Therefore, with $\tilde{\phi}_1 = \frac{1}{\|Q\|_{L^2}} Q$, we have $L_- \tilde{\phi}_1 = 0$.

Now, we shall show that there exists a double eigenvalue $\tilde{e}_2 = \tilde{e}_3 = -2\epsilon^2/3 + O(\epsilon^3)$ of L_- . In other words, we need to solve $L_- \tilde{\phi} = e \tilde{\phi}$ for e close to $e_1 - E = O(\epsilon^2)$. Again, we write $\tilde{\phi} = z \cdot \phi + g$, where g is in $\mathbf{V}^\perp$ and $z \in \mathbb{R}^3$. Then

$$(H_0 - e_1)g = (e + \lambda(\epsilon^2 - Q^2))(z \cdot \phi + g). \quad (4.8)$$

This equation is solvable if and only if

$$(\phi_j, (e + \lambda\epsilon^2)z \cdot \phi - \lambda Q^2 z \cdot \phi) = \lambda(\phi_j, Q^2 g) \quad \forall j = 1, 2, 3.$$

Since $Q^2 = \rho^2 \phi_1^2 + 2\rho \phi_1 h + h^2$ where h is a function of $x_1, |x'|$ which is odd in x_1 , we get

$$\{e + \lambda\epsilon^2 - \lambda[\rho^2(\phi_1^2, \phi_j^2) + 2\rho(\phi_j^2 \phi_1, h) + (\phi_j^2, h^2)]\} z_j = \lambda(\phi_j, Q^2 g). \quad (4.9)$$

Moreover, since $Q_E(x)$ depends on $x_1, |x'|$ and it is odd in x_1 for $x = (x_1, x')$, we see that when we restrict g to the class of function which is odd in x_2 and even in x_1, x_3 we have $(\phi_j, Q^2 g) = 0$ for $j = 1, 3$. Solving the equation (4.9) in this class of functions g , we obtain an eigenvalue

$$e = \tilde{e}_2 := \lambda [I\rho^2 + 2\rho(\phi_2^2 \phi_1, h) + (\phi_2^2, h^2) - \epsilon^2] + O(\epsilon^4) = -\frac{2}{3}\lambda\epsilon^2 + O(\epsilon^3)$$

of L_- with a normalized eigenfunction $\tilde{\phi}_2 = \phi_2 + g_2$, where $g_2 = O(\epsilon^2)$ which is odd in x_2 and even in x_1, x_3 .

Now, let $\tilde{\phi}_3 = (R_{23}(\pi/2), 1) * \tilde{\phi}_2$. Since $(R_{23}(\pi/2), 1) * Q_E = Q_E$, we see that $L_- \tilde{\phi}_3 = \tilde{e}_2 \tilde{\phi}_3$. Moreover, $\tilde{\phi}_3$ is odd in x_3 and even in x_1, x_2 . Therefore, $(\tilde{\phi}_2, \tilde{\phi}_3) = 0$. So, $\tilde{e}_2$ has multiplicity at least two. Then, by counting the number of discrete eigenvalues of L_- and H , we see that $\tilde{e}_2$ has multiplicity two and we have found all of the discrete eigenvalues of L_- . This completes the proof of the lemma. $\square$

Lemma 4.5 (Spectrum of L_+). *For $Q = Q_E$, the following statements hold*

- (i) *The discrete spectrum of L_+ consists of $\{\hat{e}_0, 0, \hat{e}_1\}$ where $\hat{e}_0 = e_0 - e_1 + O(\epsilon^2)$ and $\hat{e}_1 = 2\lambda\epsilon^2 + O(\epsilon^3)$ are simple eigenvalues, and 0 is a double eigenvalue. For $j = 0, 1, 2, 3$, there exist localized orthonormal functions $\varphi_j = \phi_j + O(\epsilon^2)$ such that*

$$L_+ \varphi_k = \hat{e}_k \varphi_k, \quad \text{for } k = 0, 1, \text{ and } L_+ \varphi_j = 0, \text{ for } j = 2, 3.$$

Moreover, $\varphi_0(x)$ is even with respect to x_2, x_3 and $\varphi_j(x)$ are odd with respect to x_j and even with respect to x_k for all $j, k > 1$ and $j \neq k$.

- (ii) *The continuous spectrum of L_+ is $\sigma_c(L_+) = [|E|, \infty)$.*

Proof. Again, we only need to prove (i). The existence of φ_0 and $\hat{e}_0$ follows from standard perturbation theory. Now consider the eigenvalues which are a perturbation of $e_1 - E$. Recall that $Q_E(\Gamma_0(r_2, r_3)x)$ are all solutions, where $r_2, r_3 \in [0, \pi)^2$. From (2.2) and (2.3), we have

$$\Gamma_0 = \Gamma_0(r_2, r_3) = \begin{bmatrix} \cos(r_2) \cos(r_3) & -\sin(r_2) & \cos(r_2) \sin(r_3) \\ \sin(r_2) \cos(r_3) & \cos(r_2) & \sin(r_2) \sin(r_3) \\ \sin(r_3) & 0 & \cos(r_3) \end{bmatrix}. \quad (4.10)$$

In particular,

$$\frac{\partial \Gamma_0(0, 0)}{\partial r_2} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \frac{\partial \Gamma_0(0, 0)}{\partial r_3} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}. \quad (4.11)$$

Now, let

$$Z_1 := \frac{\partial Q_E}{\partial E}, \quad Z_2 := \frac{\partial Q_E(\Gamma_0(r_2, r_3)x)}{\partial r_2} \Big|_{r_2=r_3=0}, \quad Z_3 := \frac{\partial Q_E(\Gamma_0(r_2, r_3)x)}{\partial r_2} \Big|_{r_2=r_3=0}. \quad (4.12)$$

From (4.11), we see that

$$\begin{aligned} Z_1(x) &= \partial_E[\rho(\epsilon)]\phi_1 + \partial_E h, \\ Z_2(x) &= (x_2\partial_1 - x_1\partial_2)Q_E(x) = \rho(\epsilon)\phi_2 + (x_2\partial_1 - x_1\partial_2)h, \\ Z_3(x) &= (x_3\partial_1 - x_1\partial_3)Q_E(x) = \rho(\epsilon)\phi_3 + (x_3\partial_1 - x_1\partial_3)h. \end{aligned}$$

Since $Q_E(x)$ is even with respect to x_2 and x_3 , so is Z_1 . Moreover, Z_2 is odd with respect to x_2 and even with respect to x_3 and Z_3 is odd with respect to x_3 and even with respect to x_2 . Therefore, we have

$$(Z_1, Z_2) = (Z_1, Z_3) = (Z_2, Z_3) = 0. \quad (4.13)$$

Now, recall the equation (3.1) for the excited states:

$$(H_0 - E)Q + \lambda|Q|^2Q = 0.$$

Taking $Q = Q_E(\Gamma_0(r_2, r_3)x)$ and differentiating this equation with respect to E , r_2 and r_3 , we find

$$L_+ Z_1 = Q_E, \quad L_+ Z_2 = L_+ Z_3 = 0.$$

So

$$\varphi_2 = \frac{1}{\|Z_2\|_{L^2}} Z_2, \quad \varphi_3 = \frac{1}{\|Z_3\|_{L^2}} Z_3$$

are (orthonormal) zero-eigenfunctions. The computation of φ_1 and $\hat{e}_1$ is exactly the same as in Lemma 4.4. In particular, as in (4.9), we get

$$\hat{e}_1 = 3\lambda[3I\rho^2 + 2\rho(\phi_1^3, h) + (\phi_1^2, h^2)] - \lambda\epsilon^2 + O(\epsilon^4) = 2\lambda\epsilon^2 + O(\epsilon^3).$$

This completes the proof of the lemma. $\square$

Now, with Z_j as in (4.12), we have

$$\mathbf{L}_0 \begin{bmatrix} 0 \\ Q_E \end{bmatrix} = 0, \quad \mathbf{L}_0 \begin{bmatrix} Z_1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ Q_E \end{bmatrix}, \quad \mathbf{L}_0 \begin{bmatrix} Z_2 \\ 0 \end{bmatrix} = \mathbf{L}_0 \begin{bmatrix} Z_3 \\ 0 \end{bmatrix} = 0. \quad (4.14)$$

Moreover, from Lemma 4.4, we have $\text{kernel}(L_-) = \text{span}\{Q_E\}$. Since $(Q_E, Z_2) = (Q_E, Z_3) = 0$, there are two functions Y_2 and Y_3 such that

$$L_- Y_2 = Z_2 \quad L_- Y_3 = Z_3 \quad \text{and} \quad (Q_E, Y_2) = (Q_E, Y_3) = (Y_2, Y_3) = 0. \quad (4.15)$$

Lemma 4.6. *For all $j = 1, 2, 3$ let $\alpha_j := -(Z_j, Y_j)^{-1}$ where $Y_1 = Q_E$. Then, α_j is finite for all $j = 1, 2, 3$.*

Proof. We need to show that $(Z_j, Y_j) \neq 0$ for all $j = 1, 2, 3$. For $j = 1$, we have $(Y_1, Z_1) = (Q_E, \partial_E Q_E) = (6I)^{-1} + O(\epsilon) \neq 0$ for ϵ sufficiently small. Now, we shall show that $(Z_j, Y_j) \neq 0$ for ϵ sufficiently small. Since the proof for the case $j = 2$ is identical to that of the case $j = 3$, we only need to prove $(Z_2, Y_2) \neq 0$. Recall that $Z_2 = a(\epsilon)\phi_2 + \tilde{h}$ for some localized order ϵ^3 function $\tilde{h}$ which is odd in x_2 and even in x_3 . We write

$$Y_2 = \sum_{j=0}^3 b_j \tilde{\phi}_j + g, \quad g \perp \tilde{\phi}_j, \quad \forall j = 0, 1, 2, 3.$$

Here $\tilde{\phi}_j$ are the eigenfunctions of L_- defined in Lemma 4.4. By choosing Y_2 not to have a component in the kernel of L_- , we can assume $b_1 = 0$. Moreover, since $L_- Y_2 = Z_2$ we have

$$L_- g = - \sum_{j=0}^3 \tilde{e}_j \tilde{\phi}_j + \rho(\epsilon)\phi_2 + \tilde{h}.$$

Taking the inner product of this equation with $\tilde{\phi}_j$ for $j = 0, 1, 2, 3$ and using the symmetry properties of these functions, we get

$$\tilde{e}_2 b_2 = \rho(\epsilon)(\tilde{\phi}_2, \phi_2) + (\tilde{\phi}_2, \tilde{h}) = \rho(\epsilon) + O(\epsilon^3), \quad b_j = 0, \quad \forall j \neq 2.$$

Since $\tilde{e}_2 = -2\lambda\epsilon^2/3 + O(\epsilon^3)$, $\tilde{\phi}_2 = \phi_2 + O(\epsilon^2)$ we see that

$$b_2 = -\frac{3\lambda\epsilon^{-1}}{2(3I)^{1/2}} + O(1).$$

To sum up, we have $Y_2 = (\epsilon^{-1})\tilde{\phi}_2 + g$ for $g \in \mathbf{P}_c(L_-)$ and satisfies

$$L_- g = O(\epsilon^3)\phi_2 + O(\epsilon^3) = O(\epsilon^3).$$

This implies that $g = O(\epsilon)$. Therefore $(Y_2, Z_2) = O(1) + O(\epsilon) \neq 0$ for ϵ sufficiently small. This completes the proof of the Lemma. $\square$

Now, we have the following theorem on the spectrum of $\mathbf{L}_0$:

Theorem 4.7 (Invariant Subspaces of $\mathbf{L}_0$ - Resonant Case). *Assume that $e_0 < 2e_1$. The space $\mathbf{E} := L^2(\mathbb{R}^3, \mathbb{C}^2)$ can be decomposed as the direct sum of $\mathbf{L}_0$ -invariant subspaces*

$$\mathbf{E} = \oplus_{j=1}^3 \mathbf{E}_{0j} \oplus \mathbf{E}_+ \oplus \mathbf{E}_- \oplus \mathbf{E}_c. \quad (4.16)$$

If f and g belong to different subspaces, then $\langle Jf, g \rangle = 0$. These subspaces and their corresponding projections satisfy the following:

(i) For each $j = 1, 2, 3$, let

$$\Phi_{0j} := \frac{\partial}{\partial r_j} \overrightarrow{e^{-ir_1} Q_E(\Gamma_0(r_2, r_3)x)} \Big|_{r_1=r_2=r_3=0}.$$

Also, let $\tilde{\Phi}_{01} = \frac{\partial Q_E}{\partial E}$ and $\tilde{\Phi}_{0k} := \begin{bmatrix} 0 \\ Y_k \end{bmatrix}$ for $k = 2, 3$. Then, we have

$\mathbf{L}_0 \tilde{\Phi}_{0j} = \Phi_{0j}$, $\mathbf{L}_0 \Phi_{0j} = 0$ and

$$\alpha_j^{-1} = \langle J\Phi_{0j}, \tilde{\Phi}_{0j} \rangle, \quad \langle J\Phi_{0j}, \Phi_{0j} \rangle = \langle J\tilde{\Phi}_{0j}, \tilde{\Phi}_{0j} \rangle = 0.$$

Moreover, the subspace $\mathbf{E}_{0j}$ is spanned by $\{\Phi_{0j}, \tilde{\Phi}_{0j}\}$ with the projection $\mathbf{P}_{0j}$ from $\mathbf{E}$ onto $\mathbf{E}_{0j}$ defined by

$$\mathbf{P}_{0j}f = -\alpha_j \langle J\tilde{\Phi}_{0j}, f \rangle \Phi_{0j} + \alpha_j \langle J\Phi_{0j}, f \rangle \tilde{\Phi}_{0j}, \quad \forall f \in \mathbf{E}, \quad j = 1, 2, 3.$$

(ii) There are four single eigenvalues $\omega_1, \omega_2 := -\bar{\omega}_1, \omega_3 := \bar{\omega}_1, \omega_4 := -\omega_1$ of $\mathbf{L}_0$ with corresponding eigenvectors $\Phi_1, \Phi_2 := \sigma_3 \bar{\Phi}_1, \Phi_3 := \bar{\Phi}_1, \Phi_4 := \sigma_3 \Phi_1$ where $\Phi_1 := \begin{bmatrix} \phi_0 \\ -i\phi_0 \end{bmatrix} + h_1$, $\omega_1 := i\kappa + \gamma$ with $\kappa := e_1 - e_0 + O(\epsilon^2) > 0$ and $\gamma := \gamma_0 \epsilon^4 + O(\epsilon^5)$ for some positive order one constant γ_0 . The function h_1 satisfies $\|h_1\|_{L^p} \leq C[\epsilon^2 + \epsilon^{6-\frac{12}{p}}]$ for all $1 \leq p \leq \infty$ and $\|h_1\|_{L_{\text{loc}}^2} \leq C\epsilon^2$ and $(J\Phi_1, \Phi_1) = 0$. The subspaces $\mathbf{E}_+$, $\mathbf{E}_-$ are defined as

$$\mathbf{E}_+ := \text{span}_{\mathbb{C}} \{\Phi_1, \Phi_2\}, \quad \mathbf{E}_- := \text{span}_{\mathbb{C}} \{\Phi_3, \Phi_4\}.$$

The projections of $\mathbf{E}$ onto $\mathbf{E}_+$ and $\mathbf{E}_-$ are defined by

$$\begin{aligned} \mathbf{P}_+ f &= c_1 \langle J\Phi_1, f \rangle \Phi_2 - \bar{c}_1 \langle J\Phi_2, f \rangle \Phi_1, \\ \mathbf{P}_- f &= \bar{c}_1 \langle J\Phi_3, f \rangle \Phi_4 - c_1 \langle J\Phi_4, f \rangle \Phi_3, \quad \text{with } c_1 := \langle J\Phi_1, \Phi_2 \rangle^{-1}. \end{aligned}$$

(iii) $\mathbf{E}_c = \{g : \langle Jf, g \rangle = 0, \quad \forall f \in \oplus_{j=0}^3 \mathbf{E}_{0j} \oplus \mathbf{E}_+ \oplus \mathbf{E}_-\}$. Its corresponding projection is $\mathbf{P}_c(\mathbf{L}_0) = Id - \sum_{j=1}^3 \mathbf{P}_{0j}(\mathbf{L}_0) - \mathbf{P}_1^+(\mathbf{L}_0) - \mathbf{P}_1^-(\mathbf{L}_0)$.




Figure 1: Spectrum of $\mathbf{L}_0$ in the resonant case.

Proof. From (4.14), we have $\mathbf{L}_0 \Phi_{0j} = 0$ and $\mathbf{L}_0 \tilde{\Phi}_{0j} = \Phi_{0j}$ for all $k = 1, 2, 3$. So, from (4.7), (4.14), (4.15) and Lemma 4.6, the statement (i) follows. Moreover, we have

$$\mathbf{L}_0|_{\mathbf{E}_0} = \begin{bmatrix} 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The proofs of (ii) and (iii) are similar and simpler than the proofs of (iii) and (iv) of Theorem 4.11 below, so they are omitted. $\square$

Now, if we denote

$$\mathbf{L}_r := \text{the linearized operator around excited state } e^{ir_1} Q_E(\Gamma_0(r_2, r_3)x),$$

for $r = (r_1, r_2, r_3) \in \mathbb{R}^3$, then

$$\mathbf{L}_r \left(r * \begin{bmatrix} u \\ v \end{bmatrix} \right) = r * \left(\mathbf{L}_0 \begin{bmatrix} u \\ v \end{bmatrix} \right), \text{ where } r * \begin{bmatrix} f(x) \\ g(x) \end{bmatrix} := e^{Jr_1} \begin{bmatrix} f(\Gamma_0(r_2, r_3)x) \\ g(\Gamma_0(r_2, r_3)x) \end{bmatrix}.$$

This observation leads to the following corollary.

Corollary 4.8. For $r = (r_1, r_2, r_3) \in \mathbb{R}^3$ and for $j = 1, 2, 3$, let $\Phi_{0j}^r := r * \Phi_{0j}$ and $\tilde{\Phi}_{0j}^r = r * \tilde{\Phi}_{0j}$. Similarly, for $j = 1, 2, 3, 4$, let $\Phi_j^r := r * \Phi_j$. Also, let

$$\begin{aligned}\mathbf{E}_{j0}^r &:= \text{span}\{\Phi_{0j}^r, \tilde{\Phi}_{0j}^r\} \quad j = 1, 2, 3, \\ \mathbf{E}_+^r &:= \text{span}\{\Phi_1^r, \Phi_2^r\}, \quad \mathbf{E}_-^r := \text{span}\{\Phi_3^r, \Phi_4^r\} \quad \forall j = 1, 2, 3, \\ \mathbf{E}_c^r &:= \{f \in \mathbf{E} : \langle Jf, g \rangle = 0, \quad \forall g \in \oplus_{j=1}^3 \mathbf{E}_{0j}^r \oplus \mathbf{E}_-^r \oplus \mathbf{E}_+^r\}.\end{aligned}$$

Then $\mathbf{E}_{0j}^r, \mathbf{E}_k^r$ are invariant under $\mathbf{L}_r$, $\mathbf{E} = \oplus_{j=1}^3 \mathbf{E}_{0j}^r \oplus \mathbf{E}_-^r \oplus \mathbf{E}_+^r \oplus \mathbf{E}_c^r$, and the projections from $\mathbf{E}$ into these subspaces are defined exactly as the corresponding projections in Theorem 4.7. Moreover,

$$\mathbf{L}_r \tilde{\Phi}_{0j}^r = \Phi_{0j}^r, \quad \mathbf{L}_r \Phi_{0j}^r = 0, \quad \mathbf{L}_r \Phi_k^r = \omega_k \Phi_k^r, \quad \forall j = 1, 2, 3, \quad k = 1, 2, 3, 4.$$

Analogous to Theorem 4.7, we have the following for the non-resonant case:

Theorem 4.9 (Invariant Subspaces of $\mathbf{L}_0$ – Non-Resonant Case). Assume that $e_0 > 2e_1$. Then, the space $\mathbf{E} := L^2(\mathbb{R}^3, \mathbb{C}^2)$ can be decomposed as the direct sum of $\mathbf{L}_0$ -invariant subspaces

$$\mathbf{E} = \oplus_{j=1}^3 \mathbf{E}_{0j} \oplus \mathbf{E}_1 \oplus \mathbf{E}_2 \oplus \mathbf{E}_c. \quad (4.17)$$

If f and g belong to different subspaces, then $\langle Jf, g \rangle = 0$. These subspaces and their corresponding projections satisfy the followings

(i) For each $j = 1, 2, 3$, let

$$\Phi_{0j} := \frac{\partial}{\partial r_j} \overrightarrow{e^{-ir_1} Q_E(\Gamma_0(r_2, r_3)x)}|_{r_1=r_2=r_3=0}.$$

Also, let $\tilde{\Phi}_{01} = \frac{\partial \overrightarrow{Q_E}}{\partial E}$ and $\tilde{\Phi}_{0k} := \begin{bmatrix} 0 \\ Y_k \end{bmatrix}$ for $k = 2, 3$. Then, we have

$$\mathbf{L}_0 \tilde{\Phi}_{0j} = \Phi_{0j}, \quad \mathbf{L}_0 \Phi_{0j} = 0 \quad \text{and}$$

$$\alpha_j^{-1} = \langle J\Phi_{0j}, \tilde{\Phi}_{0j} \rangle, \quad \langle J\Phi_{0j}, \Phi_{0j} \rangle = \langle J\tilde{\Phi}_{0j}, \tilde{\Phi}_{0j} \rangle = 0.$$

Moreover, the subspace $\mathbf{E}_{0j}$ is spanned by $\{\Phi_{0j}, \tilde{\Phi}_{0j}\}$ with the projection $\mathbf{P}_{0j}$ from $\mathbf{E}$ onto $\mathbf{E}_{0j}$ defined by

$$\mathbf{P}_{0j} f = -\alpha_j \langle J\tilde{\Phi}_{0j}, f \rangle \Phi_{0j} + \alpha_j \langle J\Phi_{0j}, f \rangle \tilde{\Phi}_{0j}, \quad \forall f \in \mathbf{E}, \quad j = 1, 2, 3.$$

(ii) There are two single purely imaginary eigenvalues $\omega_1, \omega_2 := -\omega_1$ of $\mathbf{L}_0$ with eigenvectors Φ_1 and $\Phi_2 := \bar{\Phi}_1$ where $\Phi_1 = \begin{bmatrix} u \\ -iv \end{bmatrix}$, $\omega_1 = i\kappa$ for some

constant $\kappa = e_1 - e_0 + O(\epsilon^2) > 0$. The functions u and v are real with $(J\Phi_1, \Phi_1) = 2i(u, v) = i$ and satisfy $L_+u = -\kappa v, L_-v = -\kappa u$. For each $j = 1, 2$, the subspace $\mathbf{E}_j$ is spanned by Φ_j , and the projection of $\mathbf{E}$ onto $\mathbf{E}_j$ is defined by

$$\mathbf{P}_j(\mathbf{L}_0)f = (-1)^j i \langle J\Phi_j, f \rangle \Phi_j, \quad \forall f \in \mathbf{E}.$$

(iii) $\mathbf{E}_c = \{g : \langle Jf, g \rangle = 0, \quad \forall f \in \oplus_{j=1}^3 \mathbf{E}_{0j} \oplus \mathbf{E}_1 \oplus \mathbf{E}_2\}$. Its corresponding projection is $\mathbf{P}_c(\mathbf{L}_0) = Id - \sum_{j=1}^3 \mathbf{P}_{0j}(\mathbf{L}_0) - \sum_{j=1}^2 \mathbf{P}_j(\mathbf{L}_0)$.




Figure 2: Spectrum of $\mathbf{L}_0$ in the non-resonant case.

Corollary 4.10. For $r = (r_1, r_2, r_3) \in \mathbb{R}^3$ and for $j = 1, 2, 3$, let $\Phi_{0j}^r := r * \Phi_{0j}$ and $\tilde{\Phi}_{0j}^r := r * \tilde{\Phi}_{0j}$. Similarly, for $j = 1, 2$, let $\Phi_j^r := r * \Phi_j$. Also, let

$$\mathbf{E}_{j0}^r := \text{span}\{\Phi_{0j}^r, \tilde{\Phi}_{0j}^r\} \quad j = 1, 2, 3,$$

$$\mathbf{E}_k^r := \text{span}\{\Phi_k^r\}, \quad \forall k = 1, 2,$$

$$\mathbf{E}_c^r := \{f \in \mathbf{E} : \langle Jf, g \rangle = 0, \quad \forall g \in \oplus_{j=1}^3 \mathbf{E}_{0j}^r \oplus \mathbf{E}_1^r \oplus \mathbf{E}_2^r\}$$

Then $\mathbf{E}_{0j}^r, \mathbf{E}_k^r$ are invariant under $\mathbf{L}_r$, $\mathbf{E} = \oplus_{j=1}^3 \mathbf{E}_{0j}^r \oplus \mathbf{E}_1^r \oplus \mathbf{E}_2^r \oplus \mathbf{E}_c^r$, and the projections from $\mathbf{E}$ into these subspaces are defined exactly as those corresponding projections in Theorem 4.9. Moreover,

$$\mathbf{L}_r \tilde{\Phi}_{0j}^r = \Phi_{0j}^r, \quad \mathbf{L}_r \Phi_{0j}^r = 0, \quad \mathbf{L}_r \Phi_k^r = \omega_k \Phi_k^r, \quad \forall j = 1, 2, 3, \quad k = 1, 2.$$

4.2 Spectral Analysis of $\mathbf{H}_0$

Recall (3.17) that $\tilde{Q}_E = \epsilon[v_2 + h_2(\epsilon) + h(\epsilon, v_2)]$ where $v_2 = 1/(2\sqrt{I})[\phi_1 + i\phi_2]$, $h_2(\epsilon) = O(\epsilon)v_2$ and $h(\epsilon, v_2) \in \mathbf{V}^\perp$. In other word, we can write $\tilde{Q}_E = \tilde{\rho}(\epsilon)[\phi_1 + i\phi_2] + \tilde{h}$ where $\tilde{\rho}(\epsilon) = \epsilon/(2\sqrt{I}) + O(\epsilon^2)$ and $\tilde{h} \in \mathbf{V}^\perp$ and of order $O(\epsilon^3)$. From (4.2), the linearized operator around $\tilde{Q}_E$ is

$$\mathbf{H}_0 = J(H_0 - E) + W_{\tilde{Q}_E}, \quad W_{\tilde{Q}_E} := \begin{bmatrix} W_{1, \tilde{Q}_E} & W_{2, \tilde{Q}_E} \\ W_{3, \tilde{Q}_E} & W_{4, \tilde{Q}_E} \end{bmatrix}, \quad (4.18)$$

where

$$\begin{aligned} W_{1,Q} &= 2\lambda \operatorname{Re} Q \operatorname{Im} Q, & W_{2,Q} &= \lambda[(\operatorname{Re} Q)^2 + 3(\operatorname{Im} Q)^2], \\ W_{3,Q} &= -\lambda[3(\operatorname{Re} Q)^2 + (\operatorname{Im} Q)^2], & W_{4,Q} &= -W_{1,Q}. \end{aligned} \quad (4.19)$$

Now let

$$\mathbf{K}_0 := \begin{bmatrix} H_0 - E - W_3 & W_1 \\ W_1 & H_0 - E + W_2 \end{bmatrix}. \quad (4.20)$$

We see that $\mathbf{K}_0$ is self adjoint and

$$\mathbf{H}_0 = J\mathbf{K}_0, \quad \mathbf{H}_0^* = -\mathbf{K}_0 J.$$

Moreover, denoting

$$\mathbf{H}_r = \text{linearized operator around excited state } e^{ir_1} \tilde{Q}_E(\Gamma_1(r_2, r_3)x)$$

for $r = (r_1, r_2, r_3) \in \mathbb{R}^3$, we have

$$\mathbf{H}_r \left(r * \begin{bmatrix} u \\ v \end{bmatrix} \right) = r * \left(\mathbf{H}_0 \begin{bmatrix} u \\ v \end{bmatrix} \right), \quad \text{where } r * \begin{bmatrix} f(x) \\ g(x) \end{bmatrix} = e^{Jr_1} \begin{bmatrix} f(\Gamma_1(r_2, r_3)x) \\ g(\Gamma_1(r_2, r_3)x) \end{bmatrix}. \quad (4.21)$$

Therefore, as above, to study the spectrum of $\mathbf{H}_r$, it suffices to study the spectrum of $\mathbf{H}_0$. We have the following theorem:

Theorem 4.11 ($\mathbf{H}_0$ -Invariant Subspaces – Resonant Case). *Assume that $e_0 < 2e_1$. The space $\mathbf{E} = L^2(\mathbb{R}^3, \mathbb{C}^2)$ can be decomposed into $\mathbf{H}_0$ -invariant subspaces as*

$$\mathbf{E} = \mathbf{E}_{01} \oplus \mathbf{E}_{02} \oplus \mathbf{E}_1 \oplus \mathbf{E}_2 \oplus \mathbf{E}_+ \oplus \mathbf{E}_- \oplus \mathbf{E}_c.$$

If f and g belong to different subspaces, then $\langle Jf, g \rangle = 0$. These subspaces and their corresponding projections satisfy the following:

(i) *The subspaces*

$$\mathbf{E}_{01} := \text{span}_{\mathbb{C}} \left\{ \Phi_{00} := \frac{\partial}{\partial_E} \overrightarrow{\widetilde{Q}_E}, \Phi_{01} := \frac{\partial}{\partial r_1} \overrightarrow{e^{-ir_1} \widetilde{Q}_E} \Big|_{r_1=0} \right\}$$

and

$$\mathbf{E}_{02} := \text{span}_{\mathbb{C}} \left\{ \Phi_{0j} := \frac{\partial}{\partial r_j} \overrightarrow{\widetilde{Q}_E}(\Gamma_1(r_2, r_3)x) \Big|_{r_2=r_3=0} \quad j = 2, 3 \right\}.$$

Moreover, $\mathbf{H}_0 \Phi_{00} = \Phi_{01}$, $\mathbf{H}_0 \Phi_{0j} = 0$, $\forall j = 1, 2, 3$ and

$$\langle J\Phi_{0j}, \Phi_{0k} \rangle = 0, \quad \text{if } (j, k) \notin \{(0, 1), (1, 0), (2, 3), (3, 2)\}.$$

The projections $\mathbf{P}_0(\mathbf{H}_0) : \mathbf{E} \rightarrow \mathbf{E}_0 := \mathbf{E}_{01} \oplus \mathbf{E}_{02}$ are defined by $\mathbf{P}_0(\mathbf{H}_0) = \mathbf{P}_{01}(\mathbf{H}_0) + \mathbf{P}_{02}(\mathbf{H}_0)$ with

$$\begin{aligned} \mathbf{P}_{01}(\mathbf{H}_0)f &= \beta_1 \langle J\Phi_{00}, f \rangle \Phi_{01} - \beta_1 \langle J\Phi_{01}, f \rangle \Phi_{00}, \\ \mathbf{P}_{02}(\mathbf{H}_0)f &= \beta_2 \langle J\Phi_{02}, f \rangle \Phi_{03} - \beta_2 \langle J\Phi_{03}, f \rangle \Phi_{02}, \quad \forall f \in \mathbf{E}, \end{aligned}$$

where $\beta_1 := \langle J\Phi_{00}, \Phi_{01} \rangle^{-1} = O(1)$ and $\beta_2 := \langle J\Phi_{02}, \Phi_{03} \rangle^{-1} = O(\epsilon^{-2})$.

(ii) *There exist $\Phi_1 := \begin{bmatrix} -i\phi_1 + \phi_2 \\ \phi_1 + i\phi_2 \end{bmatrix} + O(\epsilon^2)$, $\Phi_2 := \bar{\Phi}_1$ and purely imaginary numbers $\omega_1 := i(\lambda\epsilon^2 + O(\epsilon^3))$ and $\omega_2 := \bar{\omega}_1$ such that for $j = 1, 2$, the subspace $\mathbf{E}_j$ is spanned by Φ_j and*

$$\mathbf{H}_0 \Phi_j = \omega_j \Phi_j, \quad \langle J\Phi_1, \Phi_1 \rangle = -4i, \quad \langle J\Phi_1, \Phi_2 \rangle = 0.$$

The projection $\mathbf{P}_j(\mathbf{H}_0) : \mathbf{E} \rightarrow \mathbf{E}_j$ is defined by

$$\mathbf{P}_1(\mathbf{H}_0)f = -\frac{1}{4i} \langle J\Phi_1, f \rangle \Phi_1, \quad \mathbf{P}_2(\mathbf{H}_0)f = \frac{1}{4i} \langle J\Phi_2, f \rangle \Phi_2, \quad \forall f \in \mathbf{E}.$$

(iii) *For $j = 3, 4, 5, 6$ there exist eigenfunctions Φ_j with corresponding eigenvalues ω_j satisfying the condition $\omega_{k+1} = \bar{\omega}_k$, $\Phi_{k+1} = \bar{\Phi}_k$ for $k = 3, 5$, and $\text{Im}\omega_4 = \kappa = e_1 - e_0 + O(\epsilon^2) > 0$, $\text{Re}\omega_4 = \gamma = \gamma_0\epsilon^4 + O(\epsilon^6)$ for some positive order one constant γ_0 . Moreover, we have*

$$\omega_5 = -\omega_3, \quad \langle J\Phi_5, \Phi_4 \rangle = 1, \quad \langle J\Phi_j, \Phi_j \rangle = 0 \quad \text{for all } j = 3, 4, 5, 6.$$

The subspaces $\mathbf{E}_+ = \text{span}\{\Phi_4, \Phi_5\}$ and $\mathbf{E}_- = \text{span}\{\Phi_3, \Phi_6\}$ with the projection $\mathbf{P}_{\pm}(\mathbf{H}_0) : \mathbf{E} \rightarrow \mathbf{E}_{\pm}$ defined by

$$\begin{aligned} \mathbf{P}_+(\mathbf{H}_0)f &= \langle J\Phi_5, f \rangle \Phi_4 - \langle J\Phi_4, f \rangle \Phi_5, \\ \mathbf{P}_-(\mathbf{H}_0)f &= \langle J\Phi_6, f \rangle \Phi_3 - \langle J\Phi_3, f \rangle \Phi_6, \quad \forall f \in \mathbf{E}. \end{aligned}$$

- (iv) $\mathbf{E}_c = \{g \in \mathbf{E} : \langle Jf, g \rangle = 0, \quad \forall f \in \mathbf{E}_0 \oplus \mathbf{E}_1 \oplus \mathbf{E}_2 \oplus \mathbf{E}_+ \oplus \mathbf{E}_-\}$. Its corresponding projection is $\mathbf{P}_c(\mathbf{H}_0) = \text{Id} - \sum_{j=0}^2 \mathbf{P}_j(\mathbf{H}_0) - \mathbf{P}_+(\mathbf{H}_0) - \mathbf{P}_-(\mathbf{H}_0)$.




Figure 3: Spectrum of $\mathbf{H}_0$ in the resonant case.

Proof. From Lemma 4.1 and Lemma 4.2, we see that $\mathbf{H}_0$ has no eigenvalues in $\Sigma_c \cup \{z : \text{dist}(z, \Sigma_p) \geq \epsilon\}$. Therefore, we shall only need to look for eigenvalues of $\mathbf{H}_0$ in $\{z : \text{dist}(z, \Sigma_p) \leq \epsilon \text{ and } z \notin \Sigma_c\}$. First of all, note that $\mathbf{H}_0$ is a small perturbation of the operator JH (recall that $H = H_0 - E$) whose discrete spectrum $\sigma_d(JH) = \{\pm i(e_0 - E), \pm i(e_1 - E)\}$, and whose eigenfunctions are $\Phi_j^\pm := \begin{bmatrix} \phi_j \\ \pm i\phi_j \end{bmatrix}$, for $j = 0, 1, 2, 3$. The eigenvalues $\pm i(e_0 - E)$ are simple and the eigenvalues $\pm i(e_1 - E) = O(\epsilon^2)$ have multiplicity 3. Moreover, the continuous spectrum of JH is $\Sigma_c := \sigma_c(J(H_0 - E)) = \{ie : |e| \geq -E\}$.

From standard perturbation theory, the dimension of eigenspaces of $\mathbf{H}_0$ for eigenvalues near 0 totals 6, and the corresponding eigenfunctions are perturbations of linear combinations of $\Phi_j^\pm, j = 1, 2, 3$. On the other hand, by the resonance condition, $|e_0 - E| = e_1 - e_0 + O(\epsilon^2) > -E$. So finding the eigenvalues of $\mathbf{H}_0$ bifurcating from $\pm i(e_1 - E)$ requires careful estimates of resolvent operators. In general, we need to solve the problem

$$\mathbf{H}_0 \begin{bmatrix} u \\ v \end{bmatrix} = \tau \begin{bmatrix} u \\ v \end{bmatrix} \quad (4.22)$$

for some functions u, v and some eigenvalue τ near zero and near $\pm i(e_1 - e_0)$.

We shall first find the eigenvalues τ near zero. Let's write $u = a \cdot \phi + h_1, v = b \cdot \phi + h_2$ where $a = (a_1, \dots, a_3)$ and $b = (b_1, \dots, b_3)$ are of order one and $h_1, h_2 \in \mathbf{V}^\perp$. Then, we have

$$\begin{cases} (H_0 - E)h_2 + W_1u + W_2v &= \tau u + (E - e_1)b \cdot \phi, \\ (H_0 - E)h_1 - W_3u - W_4v &= -\tau v + (E - e_1)a \cdot \phi. \end{cases} \quad (4.23)$$

Applying the projection $\mathbf{P}_1^\perp$, we get

$$h_1 = -(H_0 - e_1)^{-1} \mathbf{P}_1^\perp (\tau h_2 - W_3u - W_4v), \quad h_2 = (H_0 - e_1)^{-1} \mathbf{P}_1^\perp (\tau h_1 - W_1u - W_2v). \quad (4.24)$$

Taking the inner product of the equations (4.23) with ϕ_j , for $j = 1, 2, 3$, we get

$$(\phi_j, W_1u + W_2v) = \tau a_j + (E - e_1)b_j, \quad (\phi_j, W_3u + W_4v) = \tau b_j + (e_1 - E)a_j. \quad (4.25)$$

From (4.24) and (4.25), we see that $h_1, h_2, \tau = O(\epsilon^2)$. We now write $\tau = \frac{\lambda \epsilon^2}{2} \tau_1 + O(\epsilon^3)$. Then, from (4.25) we have

$$\begin{bmatrix} 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ -3 & 0 & 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ b_1 \\ b_2 \\ b_3 \end{bmatrix} = \tau_1 \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ b_1 \\ b_2 \\ b_3 \end{bmatrix}.$$

This implies that τ_1 solves the equation $\tau_1^4(\tau_1^2 + 4) = 0$. So, the eigenvalue problem (4.22) has two purely imaginary eigenvalues $\tau = \omega_1 := \lambda \epsilon^2 i + O(\epsilon^3)$ and $\tau = \omega_2 := \bar{\omega}_1$. Respectively, the eigenvectors of ω_1 and ω_2 are $\Phi_1, \bar{\Phi}_1$ with

$$\Phi_1 = \begin{bmatrix} -i\phi_1 + \phi_2 \\ \phi_1 + i\phi_2 \end{bmatrix} + O(\epsilon^2).$$

Moreover, as in the previous section, let

$$\begin{aligned} \Phi_{00} &= \frac{\partial}{\partial E} \overrightarrow{Q_E} = \partial_E \tilde{\rho}(\epsilon) \begin{bmatrix} \phi_1 \\ \phi_2 \end{bmatrix} + O(\epsilon^2), \\ \Phi_{01} &= \frac{\partial}{\partial r_1} e^{-ir_1} \overrightarrow{Q_E}|_{r_1=0} = \begin{bmatrix} \text{Im } \tilde{Q} \\ -\text{Re } \tilde{Q} \end{bmatrix} = \tilde{\rho}(\epsilon) \begin{bmatrix} \phi_2 \\ -\phi_1 \end{bmatrix} + O(\epsilon^3), \\ \Phi_{02} &= \frac{\partial}{\partial r_2} \overrightarrow{Q_E}(\Gamma_1(r_2, r_3)x)|_{r_2=r_3=0} = \tilde{\rho}(\epsilon) \begin{bmatrix} 0 \\ \phi_3 \end{bmatrix} + O(\epsilon^3), \\ \Phi_{03} &= \frac{\partial}{\partial r_3} \overrightarrow{Q_E}(\Gamma_1(r_2, r_3)x)|_{r_2=r_3=0} = \tilde{\rho}(\epsilon) \begin{bmatrix} \phi_3 \\ 0 \end{bmatrix} + O(\epsilon^3). \end{aligned} \quad (4.26)$$

We see that $\mathbf{H}_0$ has three zero - eigenvectors and one zero-generalized eigenvector. Precisely, we have

$$\mathbf{H}_0 \Phi_{00} = \Phi_{01}, \quad \mathbf{H}_0 \Phi_{0j} = 0, \quad \forall j = 1, 2, 3.$$

Next, we shall look for the eigenvalues τ of $\mathbf{H}_0$ near $\pm i(e_0 - E)$, i.e. we shall solve $(JH - \tau + W)\Phi = 0$. We shall first solve this equation for $|\tau - i(e_0 - E)| \leq \epsilon$. From Lemma 4.1, we may assume that $\text{Re } \tau \neq 0$. We write $\Phi = a\Phi_0^+ + \eta$, for some $a \in \mathbb{C}$, $\eta \perp \Phi_0^+$. Then we have

$$\begin{cases} a\tau = ai(e_0 - E) + \frac{1}{2}\langle \Phi_0^+, W(a\Phi_0^+ + \eta) \rangle, \\ (JH - \tau)\eta = -\mathbf{P}^\perp W(a\Phi_0^+ + \eta). \end{cases} \quad (4.27)$$

Here $\mathbf{P}^\perp = \mathbf{P}_d + \mathbf{P}_c$,

$$\mathbf{P}_d f := \frac{1}{2}\langle \Phi_0^-, f \rangle \Phi_0^- + \sum_{j=1}^3 \frac{1}{2}\langle \Phi_j^\pm, f \rangle \Phi_j^\pm, \quad \mathbf{P}_c f := \mathbf{P}_c(JH)f, \quad f \in \mathbf{E}.$$

Recall that $R_0(z) = (JH - z)^{-1} = R_{01}(z) + R_{02}(z)$ where,

$$R_{01}(z) := \frac{1}{2} \begin{bmatrix} i & -1 \\ 1 & i \end{bmatrix} (H - iz)^{-1}, \quad R_{02}(z) := \frac{1}{2} \begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix} (H + iz)^{-1}.$$

Therefore, $R_0(z)$ is well defined for $\text{Re } z \neq 0$. From the second equation of (4.27), we get

$$\eta = -R_0(\tau)\mathbf{P}^\perp[W(a\Phi_0^+ + \eta)]. \quad (4.28)$$

For $j = 1, 2, 3$, from (4.19) and by the symmetry properties of $\tilde{Q}$, we have

$$\begin{aligned} \langle \Phi_j^+, W\Phi_0^+ \rangle &= (\phi_0\phi_j, W_1 + iW_2) - i(\phi_0\phi_j, W_3 + iW_4) \\ &= 4i\lambda(\phi_0\phi_j, |\tilde{Q}|^2) = 0, \\ \langle \Phi_j^-, W\Phi_0^+ \rangle &= (\phi_0, \phi_j, W_1 + iW_2) + i(\phi_0\phi_j, W_3 + iW_4) \\ &= -2i\lambda(\phi_0\phi_j, \tilde{Q}^2) = 0. \end{aligned} \quad (4.29)$$

So,

$$\mathbf{P}^\perp[W\Phi_0^+] = \frac{1}{2}\langle \Phi_0^-, W\Phi_0^+ \rangle \Phi_0^- + \mathbf{P}_c W\Phi_0^+.$$

Therefore, we obtain the equation for η

$$\begin{aligned} \eta_\tau &= \frac{1}{2[i(e_0 - E) + \tau]} [a\langle \Phi_0^-, W\Phi_0^+ \rangle + \langle \Phi_0^-, W\eta \rangle] \Phi_0^- \\ &\quad - \sum_{j=1}^3 \frac{1}{2[\pm i(e_1 - E) - \tau]} \langle \Phi_j^\pm, W\eta \rangle \Phi_j^\pm - aR_0(\tau)\mathbf{P}_c W\Phi_0^+ - R_0(\tau)\mathbf{P}_c W\eta. \end{aligned} \quad (4.30)$$

Now suppose that $a = 0$. Since $\operatorname{Re}(\tau) \neq 0$ and τ is close to $i(e_0 - E)$, from (4.30) and Lemma 4.3, we see that $\|\eta\|_{L_{\text{loc}}^2} \leq C\epsilon^2 \|\eta\|_{L_{\text{loc}}^2}$. So, we get a contradiction because ϵ is small. So, without loss of generality, we may assume that $a = 1$. Then let's define $\tau_0 := i(e_0 - E) + \frac{1}{2}(\Phi_0^+, W\Phi_0^+)$. Since

$$\langle \Phi_0^+, W\Phi_0^+ \rangle = (\phi_0^2, W_1 + iW_2) - i(\phi_0^2, W_3 - iW_1) = 4i(\phi_0^2, |\tilde{Q}|^2),$$

we see that τ_0 is purely imaginary and $\operatorname{Im} \tau_0 = e_0 - e_1 + \lambda\epsilon^2 + O(\epsilon^4)$. Moreover, let

$$\begin{aligned} \tau_1 := & \frac{1}{4[i(e_0 - E) + \tau_0]} |\langle \Phi_0^-, W\Phi_0^+ \rangle|^2 - 4i\lambda^2(\phi_0|Q|^2, \frac{1}{H + i\tau_0} \mathbf{P}_c(H_0)\phi_0|Q|^2) \\ & - \lambda^2 i(\phi_0 Q^2, \frac{1}{H - i\tau_0 - i0} \mathbf{P}_c(H_0)\phi_0 Q^2). \end{aligned} \quad (4.31)$$

Then we have $|\tau_1| \lesssim \epsilon^4$. Moreover, from (1.5), and since $(H + i\tau_0)^{-1} \mathbf{P}_c$ is regular, we have $\operatorname{Re}(\tau_1) \geq \lambda^2 \lambda_0 \epsilon^4$. Here, λ and λ_0 are respectively defined in (1.1) and (1.5). From (4.27) and (4.28), we get the equation for τ

$$\tau = \tau_0 + \tau_1 - [\tau_1 + \frac{1}{2} \langle \Phi_0^+, WR_0(\tau) \mathbf{P}^\perp W\Phi_0^+ \rangle] - \frac{1}{2} \langle \Phi_0^+, WR_0(\tau) \mathbf{P}^\perp W\eta \rangle. \quad (4.32)$$

From (4.30) and Lemma 4.3 and by applying the contraction mapping theorem, for each τ in $\mathbf{D}_1$, we can find the solution η_τ in L_{-s}^2 of (4.30) such that $\|\eta_\tau\|_{L_{-s}^2} \lesssim \epsilon^2$. Moreover, for all $\tau, \tau' \in \mathbf{D}_1$, from (4.30) and Lemma 4.3, we also have

$$\|\eta_\tau - \eta_{\tau'}\|_{L_{-s}^2} \lesssim \epsilon^2 [\max\{\operatorname{Re} \tau, \operatorname{Re} \tau'\}]^{-1/2} |\tau - \tau'| + \epsilon^2 \|\eta_\tau - \eta_{\tau'}\|_{L_{-s}^2}.$$

Therefore, there exists a constant $C > 0$ such that

$$\|\eta_\tau - \eta_{\tau'}\|_{L_{-s}^2} \leq C\epsilon^2 [\max\{\operatorname{Re} \tau, \operatorname{Re} \tau'\}]^{-1/2} |\tau - \tau'|, \quad \forall \tau, \tau' \in \mathbf{D}_1. \quad (4.33)$$

Next, let

$$\mathbf{D} := \{z \in \mathbb{C} : |z - (\tau_0 + \tau_1)| \leq \epsilon^5\}. \quad (4.34)$$

Note that $\mathbf{D} \subset \mathbf{D}_1$. Also, let $f(\tau)$ be a function which is defined by the right hand side of (4.32). We shall show that f maps $\mathbf{D}$ into $\mathbf{D}$ and has a fixed point

in $\mathbf{D}$. Note that

$$\begin{aligned}
\langle \Phi_0^+, W R_0(\tau) \mathbf{P}^\perp W \Phi_0^+ \rangle &= \frac{1}{2} \langle \Phi_0^-, W \Phi_0^+ \rangle \langle \Phi_0^+, W R_0(\tau) \Phi_0^- \rangle + \langle \Phi_0^+, W R_0(\tau) \mathbf{P}_c W \Phi_0^+ \rangle \\
&= -\frac{1}{2[i(e_0 - E) + \tau]} \langle \Phi_0^-, W \Phi_0^+ \rangle \langle \Phi_0^+, W \Phi_0^- \rangle + \langle \Phi_0^+, W R_0(\tau) \mathbf{P}_c W \Phi_0^+ \rangle \\
&= -\frac{1}{2[i(e_0 - E) + \tau]} |\langle \Phi_0^-, W \Phi_0^+ \rangle|^2 + \langle \Phi_0^+, W R_0(\tau) \mathbf{P}_c W \Phi_0^+ \rangle, \\
&= -\frac{1}{2[i(e_0 - E) + \tau]} |\langle \Phi_0^-, W \Phi_0^+ \rangle|^2 + 2i\lambda^2(\phi_0 Q^2, (H - i\tau)^{-1} \mathbf{P}_c(H) \phi_0 Q^2) \\
&\quad + 4i\lambda^2(\phi_0 |Q|^2, (H + i\tau)^{-1} \mathbf{P}_c(H) \phi_0 |Q|^2).
\end{aligned} \tag{4.35}$$

Here, we have used the fact that $R_0 \mathbf{P}_c = R_{01} \mathbf{P}_c + R_{02} \mathbf{P}_c$ with

$$R_{01}(z) \mathbf{P}_c := \frac{1}{2} \begin{bmatrix} i & -1 \\ 1 & i \end{bmatrix} (H - iz)^{-2} \mathbf{P}_c(H), \quad R_{02}(z) \mathbf{P}_c := \frac{1}{2} \begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix} (H + iz)^{-1} \mathbf{P}_c(H)$$

and

$$\begin{aligned}
\langle \Phi_0^+, W R_{01}(\tau) \mathbf{P}_c W \Phi_0^+ \rangle &= 2i\lambda^2(\phi_0 Q^2, (H - i\tau)^{-1} \mathbf{P}_c(H) \phi_0 Q^2), \\
\langle \Phi_0^+, W R_{02}(\tau) \mathbf{P}_c W \Phi_0^+ \rangle &= 8i\lambda^2(\phi_0 |Q|^2, (H + i\tau)^{-1} \mathbf{P}_c(H) \phi_0 |Q|^2).
\end{aligned} \tag{4.36}$$

From (4.31), (4.35), Lemma 4.3 and $|\tau - \tau_0| \lesssim \epsilon^4$ if $\tau \in \mathbf{D}$, we obtain

$$|\tau_1 + \frac{1}{2} \langle \Phi_0^+, W R_0(\tau) \mathbf{P}^\perp W \Phi_0^+ \rangle| \lesssim \epsilon^6, \quad \forall \tau \in \mathbf{D}.$$

On the other hand,

$$\begin{aligned}
\langle \Phi_0^+, W R_0(\tau) \mathbf{P}^\perp W \eta \rangle &= -\frac{1}{2[i(e_0 - E) + \tau]} \langle \Phi_0^-, W \eta \rangle \langle \Phi_0^+, W \Phi_0^- \rangle \\
&\quad + \sum_{j=1}^3 \frac{1}{2[\pm i(e_1 - E) - \tau]} \langle \Phi_0^\pm, W \eta \rangle \langle \Phi_0^+, W \Phi_j^\pm \rangle + \langle \Phi_0^+, W R_0(\tau) \mathbf{P}_c W \eta \rangle.
\end{aligned}$$

So,

$$|\langle \Phi_0^+, W R_0(\tau) \mathbf{P}^\perp W \eta \rangle| \lesssim \epsilon^6, \quad \forall \tau \in \mathbf{D}.$$

Therefore, $|f(\tau) - (\tau_0 + \tau_1)| \lesssim \epsilon^6 \leq \epsilon^5$ for all $\tau \in \mathbf{D}$. So, $f(\tau) \in \mathbf{D}$ for all $\tau \in \mathbf{D}$. From (4.33), (4.35) and Lemma 4.3 below, we can show that there exists a constant $C > 0$ independent of ϵ such that $|f(\tau) - f(\tau')| \leq C\epsilon^2 |\tau - \tau'|$, for all τ, τ' in $\mathbf{D}$. So, the map $f : \mathbf{D} \rightarrow \mathbf{D}$ is a contraction when ϵ is sufficiently

small. Therefore, there exists unique $\tau_* \in \mathbf{D}$ such that $\tau_* = f(\tau_*)$. Moreover, we have

$$\operatorname{Im} \tau_* = \operatorname{Im} \tau_0 + O(\epsilon^2) = e_0 - e_1 + O(\epsilon^2), \quad \lambda^2 \lambda_0 \epsilon^4 \leq \operatorname{Re} \tau_* = \operatorname{Re} \tau_1 + O(\epsilon^6) \leq C\epsilon^4. \quad (4.37)$$

Next, we shall show that τ_* is the unique fixed point of f in $\mathbf{D}_1$, where $\mathbf{D}_1$ is defined in Lemma 4.3 (note that $\mathbf{D} \subset \mathbf{D}_1$). Suppose that there is $\tau' \in \mathbf{D}_1$ such that $f(\tau') = \tau'$. Using (4.33) and (4.37), we obtain

$$|\tau_* - \tau'| = |f(\tau_*) - f(\tau')| \leq C\epsilon^2 |\tau_* - \tau'| \leq \frac{1}{2} |\tau_* - \tau'|.$$

Therefore, $\tau' = \tau_*$. So, τ_* is the unique eigenvalue of $\mathbf{H}_0$ in $\mathbf{D}_1$. In summary, let $h_3 := \eta_{\tau_*}$, $\Phi_3 = \Phi_0^+ + h_3$, $\omega_3 = \tau_*$, $\Phi_4 = \bar{\Phi}_3$ and $\omega_4 = \bar{\omega}_3$. We have $\lambda_0 \lambda^2 \leq \operatorname{Re} \omega_j \lesssim \epsilon^4$, $\operatorname{Im} \omega_j = (-1)^{j-1}(e_0 - e_1) + O(\epsilon^2)$ and

$$\mathbf{H}_0 \Phi_j = \omega_j \Phi_j, \quad \forall j = 3, 4.$$

Moreover, from (4.30), we get

$$\begin{aligned} h_3 &= \frac{1}{2[i(e_0 - E + \omega_3)]} [\langle \Phi_0^-, W \Phi_0^+ \rangle + \langle \Phi_0^-, W h_3 \rangle] \Phi_0^- \\ &\quad - \sum_{j=1}^3 \frac{1}{\pm i(e_1 - E) - \omega_3} \langle \Phi_j^\pm, W h_3 \rangle \Phi_j^\pm - R_0(\omega_3) \mathbf{P}_c W \Phi_0^+ - R_0(\omega_3) \mathbf{P}_c W h_3. \end{aligned} \quad (4.38)$$

Moreover, from Lemma 4.12 below, we see that $\|h_3\|_{L^2} \leq C$. Similarly, solving the bifurcation equation $(JH + W)[\Phi_0^- + \tilde{\eta}] = z(\Phi_0^- + \tilde{\eta})$, we also obtain two other eigenvalues and eigenfunctions ω_5, ω_6 and $\Phi_5 = \Phi_0^- + h_5$, $\Phi_6 = \Phi_0^+ + h_6$ with $\omega_6 = \bar{\omega}_5$, $\Phi_6 = \bar{\Phi}_5$, $\mathbf{H}_0 \Phi_j = \omega_j \Phi_j$, $\|h_j\|_{L^2} \leq C$ for $j = 5, 6$. In particular, h_5 satisfies

$$\begin{aligned} h_5 &= \frac{1}{2[-i(e_0 - E + \omega_5)]} [\langle \Phi_0^+, W \Phi_0^- \rangle + \langle \Phi_0^+, W h_3 \rangle] \Phi_0^+ \\ &\quad - \sum_{j=1}^3 \frac{1}{\pm i(e_1 - E) - \omega_5} \langle \Phi_j^\pm, W h_3 \rangle \Phi_j^\pm - R_0(\omega_5) \mathbf{P}_c W \Phi_0^- - R_0(\omega_5) \mathbf{P}_c W h_5. \end{aligned} \quad (4.39)$$

Also,

$$\begin{aligned} \operatorname{Re} \omega_5 &= -\lambda^2 \operatorname{Im}(\phi_0 \bar{Q}^2, \frac{1}{H_0 + e_0 - 2e_1 + O(\epsilon^2) - i0} \mathbf{P}_c \phi_0 \bar{Q}^2) + O(\epsilon^6), \\ \operatorname{Im} \omega_5 &= -(e_0 - e_1) + O(\epsilon^2). \end{aligned}$$

In particular, we have $\operatorname{Re} \omega_j = O(\epsilon^4)$ and $\operatorname{Re} \omega_j \leq -\lambda_0 \lambda_0 \epsilon^4$ for $j = 5, 6$. By the uniqueness properties of these fixed points $\omega_3, \omega_4, \dots, \omega_6$, we see that they are all of eigenvalues of $\mathbf{H}_0$ in the ϵ -neighborhood of $\pm i e_{01}$. Therefore, we have found all of the eigenvalues of $\mathbf{H}_0$.

Next, we shall prove the orthogonality conditions from which the formulas of the projections follow. Firstly, from the symmetry properties of $\tilde{Q}_E$, it follows that $\langle Jf, g \rangle = 0$ if f, g are in different spaces of $\mathbf{E}_{01}$ and $\mathbf{E}_{02}$. Secondly, we claim that for any two pairs of eigenvectors, eigenvalues $(u_1, \lambda_1), (u_2, \lambda_2)$ of $\mathbf{H}$ with $\bar{\lambda}_1 + \lambda_2 \neq 0$, then $\langle Ju_1, u_2 \rangle = 0$. In fact, since $\mathbf{H} = J\mathbf{K}$ and $\mathbf{K}$ is a self-adjoint operator, we get

$$\lambda_2 \langle Ju_1, u_2 \rangle = \langle Ju_1, \mathbf{H}u_2 \rangle = \langle \mathbf{H}^* Ju_1, u_2 \rangle = \langle \mathbf{K}u_1, u_2 \rangle = -\bar{\lambda}_1 \langle Ju_1, u_2 \rangle. \quad (4.40)$$

Then, we obtain $(\bar{\lambda}_1 + \lambda_2) \langle Ju_1, u_2 \rangle = 0$. So, $\langle Ju_1, u_2 \rangle = 0$ since $\bar{\lambda}_1 + \lambda_2 \neq 0$. Therefore, $\langle Jf, g \rangle = 0$ if f and g are in different spaces of $\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_+, \mathbf{E}_-$. Moreover, $\langle J\Phi_j, \Phi_j \rangle = 0$ for all $j = 3, 4, 5, 6$. On the other hand, for $u_1 \in \mathbf{E}_0$, we have $(\mathbf{H}^*)^2 Ju_1 = (\mathbf{K}J)(\mathbf{K}J)Ju_1 = J\mathbf{H}^2 u_1 = 0$. Therefore, for all u_2 such that $\mathbf{H}u_2 = \lambda u_2$ with $\lambda \neq 0$, we get

$$\lambda^2 \langle Ju_1, u_2 \rangle = \langle Ju_1, \mathbf{H}^2 u_2 \rangle = \langle (\mathbf{H}^*)^2 u_1, u_2 \rangle = 0.$$

This proves that $\langle Jf, g \rangle = 0$ if $f \in \mathbf{E}_0$ and g is in one of $\mathbf{E}_1, \mathbf{E}_2, \mathbf{E}_+, \mathbf{E}_-$. So, we have proved all of the orthogonality conditions. Moreover, since $J\Phi_0^+ = i\Phi_0^+$ and Lemma 4.12 below, we obtain

$$\begin{aligned} \langle J\Phi_5, \Phi_4 \rangle &= \langle J(\Phi_0^+ + h_5), \Phi_0^+ + h_4 \rangle = 2i + O(\epsilon^2) + \langle Jh_5, h_4 \rangle \\ &= 2i + O(\epsilon^2) \neq 0. \end{aligned}$$

Moreover, it follows from this and (4.40) that

$$\omega_5 = -\bar{\omega}_4 = -\omega_3.$$

Finally, we shall complete the proof of Theorem 4.7 by the following two lemmas which show that all of $\omega_3, \omega_4, \dots, \omega_6$ are simple and $\|h_j\|_{L^2} \leq C$ for $j = 3, 4, \dots, 6$: $\square$

Lemma 4.12. *For $1 \leq p \leq \infty$, we have*

$$\|h_j\|_{L^p} \leq C_p[\epsilon^2 + \epsilon^{6-\frac{12}{p}}], \quad \|h_j\|_{H^1} \leq C, \quad |\langle Jh_5, h_4 \rangle| \leq C\epsilon^4, \quad \forall j = 3, 4, 5, 6.$$

Proof. To prove the first inequality of Lemma 4.12, we only need to prove it for $j = 3$. From (4.38), we only need to show that

$$\|R_0(\omega_3)\mathbf{P}_c W \Phi_0^+\|_{L^p}, \|R_0(\omega_3)\mathbf{P}_c W \eta\|_{L^p} \leq C_p[\epsilon^2 + \epsilon^{6-\frac{12}{p}}]. \quad (4.41)$$

Since $R_0(\tau) = R_{01}(\tau) + R_{02}(\tau)$ and $R_{02}(\omega_3)$ is regular and $R_{01}(\omega_3) \sim (H - i\omega_3)^{-1}$, we shall only need to prove (4.41) with $R_0(\omega_3)$ replaced by $(H - i\omega_3)^{-1}$. Now, we follow the argument in [14]. We write

$$H - i\omega_3 = H_0 - E - i\omega_3 = -\Delta - \nu^2 + V,$$

where $\nu^2 = E + i\omega_3$ and $\text{Im } \nu > 0$ and is of order ϵ^4 . By resolvent expansion, we have

$$(H - i\omega_3)^{-1}\varphi = (-\Delta - \nu^2)^{-1}\varphi + (-\Delta - \nu^2)^{-1}V(H - i\omega_3)^{-1}\varphi. \quad (4.42)$$

Because the resolvent $(-\Delta - \nu^2)^{-1}$ has the kernel $K(x) := (4\pi|x|)^{-1} \exp(i\nu|x|)$, we have

$$\begin{aligned} \|(-\Delta - \nu^2)^{-1}\varphi\|_{L^p} &\lesssim \|K * \varphi\|_{L^p} \lesssim [\|K\|_{L^p(B_1^c)} + \|K\|_{L^2(B_1)}] \|\varphi\|_{L^1 \cap L^2} \\ &\lesssim [1 + \epsilon^{4-6/p}] \|\varphi\|_{L^1 \cap L^2}. \end{aligned}$$

On the other hand, since V decays sufficiently fast, we have

$$\|V(H - i\omega_3)^{-1}\varphi\|_{L^1 \cap L^2} \lesssim \|(H - i\omega_3)^{-1}\varphi\|_{L_{\text{loc}}^2} \lesssim \|\langle x \rangle^s \varphi\|_{L^2}.$$

Then, it follows from (4.42) that

$$\|(H - i\omega_3)^{-1}\varphi\|_{L^2} \lesssim [1 + \epsilon^{4-6/p}][\|\phi\|_{L^1} + \|\langle x \rangle^s \varphi\|_{L^2}].$$

Using this estimate, we get

$$\|R_0(\omega_3)\mathbf{P}_c W \Phi_0^+\|_{L^p} + \|R_0(\omega_3)\mathbf{P}_c W \eta\|_{L^p} \leq C_p \epsilon^2 [1 + \epsilon^{4-\frac{12}{p}}].$$

So, we obtain the first inequality of Lemma 4.12. Similarly, to prove the second inequality, we only need to show that for some localized function φ

$$\|\nabla v\|_{L^2} \leq C, \quad v := (H - i\omega_3)^{-1}\varphi.$$

This follows directly by multiplying the equation $(H - i\omega_3)v = \varphi$ by $\bar{v}$ and integrating over $\mathbb{R}^3$.

Finally, we prove the last inequality of Lemma 4.41. Again, note that h_3 and h_5 are of the form

$$h_3 = \text{ok} + A_1(H - i\omega_3)^{-1}\varphi, \quad h_5 = \text{ok} + A_2(H + i\omega_5)^{-1}\varphi_*.$$

Here, $ok =$ terms which are localized and of order ϵ^2 , φ, φ_* are some localized functions of order ϵ^2 and A_1, A_2 are some constant matrices of order one. So, we get

$$\begin{aligned} |\langle Jh_5, \bar{h}_3 \rangle| &\leq C[\epsilon^4 + |((H + i\omega_5)^{-1}\varphi_*, (H + i\bar{\omega}_3)^{-1}\bar{\varphi})|] \\ &\leq C[\epsilon^4 + |((H - i\omega_3)^{-1}(H + i\omega_5)^{-1}\varphi_*, \bar{\varphi})|] \\ &\leq C\epsilon^4. \end{aligned}$$

Here, in the last step, we used the fact that for $\alpha_1, \alpha_2 \in \mathbb{C}$ such that $\text{Im } \alpha_j > 0$ and $|\text{Re } \alpha_j| \in [a_1, a_2] \subset (0, \infty)$ with $j = 1, 2$, we have

$$\|(H - \alpha_1)^{-1}(H - \alpha_2)^{-1}P_c(H)g\|_{L^2_{-s}} \leq C\|g\|_{L^2_s}, \quad \forall g \in L^2_s. \quad (4.43)$$

Here, the constant $C > 0$ is independent of α_1 and α_2 . One can prove (4.43) by using Mourre estimates and the argument as in [21], where the authors proved similar estimates for linearized operators and $\alpha_1 = \alpha_2$. For a different approach, one can see [7]. $\square$

Lemma 4.13. *The eigenvalues ω_j defined in the proof of Theorem 4.11 are simple for $j = 3, 4, 5, 6$.*

Proof. It suffices that we only prove this lemma for $j = 3$. Suppose by contradiction that there is $\tilde{\Phi}$ such that $[\mathbf{H} - \omega_3]\tilde{\Phi} = \Phi_0^+ + h_3$. Note that $h_3 = \eta = \eta(\omega_3)$. We write $\tilde{\Phi} = c\Phi_0^+ + h$ where $h \in \mathbf{E}, h \perp \Phi_0^+$ and c is in $\mathbb{C}$. Then, we have

$$c[i(e_0 - E) - \omega_3]\Phi_0^+ + W[c\Phi_0^+ + h] + (JH - \omega_3)h = \Phi_0^+ + \eta.$$

Equivalently,

$$\begin{cases} c[i(e_0 - E) + \frac{1}{2}\langle \Phi_0^+, W\Phi_0^+ \rangle - \omega_3] + \frac{1}{2}\langle \Phi_0^+, Wh \rangle = 1, \\ (JH - \omega_3)h = -c\mathbf{P}^\perp W\Phi_0^+ - \mathbf{P}^\perp Wh + \eta. \end{cases} \quad (4.44)$$

Let's now define $h^* = h - c\eta$. From (4.27) and (4.44), we see that h^* solves the equation

$$h^* = R_0(\omega_3)[\eta - \mathbf{P}^\perp Wh^*].$$

From (4.30), Lemma 4.3 and (4.43), we have

$$\|R_0(\omega_3)\eta\|_{L^2_{-s}} \leq C[\epsilon^2 + \|R_0^2(\omega_3)\mathbf{P}_c W\Phi_0^+\|_{L^2_{-s}} + \|R_0^2(\omega_3)\mathbf{P}_c W\eta\|_{L^2_{-s}}] \leq C\epsilon^2.$$

Therefore, we get $\|h^*\|_{L^2_{-s}} \leq C\epsilon^2$. Now, from the first equation of (4.44), we get

$$c[i(e_0 - E) + \frac{1}{2}\langle \Phi_0^+, W(\Phi_0^+ + \eta) \rangle - \omega_3] = 1 - \frac{1}{2}\langle \Phi_0^+, Wh^* \rangle.$$

From (4.27), we see that $i(e_0 - E) + \frac{1}{2}\langle \Phi_0^+, W[\Phi_0^+ + \eta] \rangle - \omega_3 = 0$. Therefore, we get $1 - \frac{1}{2}\langle \Phi_0^+, Wh^* \rangle = 0$. This is a contradiction since $|(\Phi_0^+, Wh^*)| \leq C\epsilon^4$. So, ω_3 is simple and the lemma follows. $\square$

As a corollary of Theorem 4.11, we obtain the same spectral properties around symmetry transformed ground states:

Corollary 4.14. *For $r = (r_1, r_2, r_3) \in \mathbb{R}^3$, and for $j = 0, 1, 2, 3$, let $\Phi_{0j}^r := r * \Phi_{0j} = e^{-Jr_1} \Phi_{0j} \circ \Gamma_1(r_2, r_3)$. Similarly, for $j = 1, 2, 3, 4, 5, 6$, let $\Phi_j^r := e^{-Jr_1} \Phi_j \circ \Gamma_1(r_2, r_3)$. Also, let $\mathbf{E}_j^r, \mathbf{E}_-^r, \mathbf{E}_+^r, \mathbf{E}_c^r$ be exactly as in Theorem 4.11, for $j = 0, 1, 2$. Then $\mathbf{E}_j^r$ are invariant under $\mathbf{H}_r$, $\mathbf{E} = \oplus_{j=0}^2 \mathbf{E}_j^r \oplus \mathbf{E}_-^r \oplus \mathbf{E}_+^r \oplus \mathbf{E}_c^r$, and the projections from $\mathbf{E}$ into these subspaces are defined exactly as those corresponding projections in Theorem 4.11. Moreover, we have*

$$\mathbf{H}_r \Phi_{00}^r = \Phi_{01}^r, \quad \mathbf{H}_r \Phi_{0j}^r = 0, \quad \mathbf{H}_r \Phi_k^r = \omega_k \Phi_k^r, \quad \forall j = 1, 2, 3, \quad k = 1, 2, 3, 4, 5, 6.$$

In the non-resonant case, we also have the following result on the spectrum of $\mathbf{H}_0$, whose proof is simpler, and is therefore skipped:

Theorem 4.15 (**$\mathbf{H}_0$ -Invariant Subspaces - Non-Resonant Case**). *Assume that $2e_1 < e_0$. Let $\mathbf{H}_0$ be defined as in (4.3). Then, the space $\mathbf{E} = L^2(\mathbb{R}^3, \mathbb{C}^2)$ can be decomposed into the $\mathbf{H}_0$ -invariant subspaces as*

$$\mathbf{E} = \oplus_{j=0}^4 \mathbf{E}_j \oplus \mathbf{E}_c.$$

If f and g belong to different subspaces, then $\langle Jf, g \rangle = 0$. These subspaces and their corresponding projections satisfy the followings:

- (i) *The subspace $\mathbf{E}_0$ is generated by zero-eigenvectors*

$$\Phi_{0j} = \frac{\partial}{\partial r_j} e^{-ir_1} \overrightarrow{\widetilde{Q}_E}(\Gamma_1(r_2, r_3)x) \Big|_{r_1=r_2=r_3=0}$$

for $j = 1, 2, 3$, and a generalized eigenvector $\Phi_{00} = \partial_E \overrightarrow{\widetilde{Q}_E}$, with $\mathbf{H}_0 \Phi_{01} = \Phi_{00}$, $\mathbf{H}_0 \Phi_{0j} = 0$, $\forall j = 0, 2, 3$ and moreover,

$$\langle J\Phi_{0j}, \Phi_{0k} \rangle = 0, \quad \text{if } (j, k) \notin \{(0, 1), (1, 0), (2, 3), (3, 2)\}.$$

The projection $\mathbf{P}_0(\mathbf{H}_0) : \mathbf{E} \rightarrow \mathbf{E}_0$ is defined by $\mathbf{P}_0(\mathbf{H}_0) = \mathbf{P}_{01}(\mathbf{H}_0) + \mathbf{P}_{02}(\mathbf{H}_0)$ with

$$\begin{aligned} \mathbf{P}_{01}(\mathbf{H}_0)f &= \beta_1 \langle J\Phi_{00}, f \rangle \Phi_{01} - \beta_1 \langle J\Phi_{01}, f \rangle \Phi_{00}, \\ \mathbf{P}_{02}(\mathbf{H}_0)f &= \beta_2 \langle J\Phi_{02}, f \rangle \Phi_{03} - \beta_2 \langle J\Phi_{03}, f \rangle \Phi_{02}, \quad \forall f \in \mathbf{E}, \end{aligned}$$

where $\beta_1 := \langle J\Phi_{00}, \Phi_{01} \rangle^{-1} = O(1)$ and $\alpha_2 = \langle J\Phi_{02}, \Phi_{03} \rangle^{-1} = O(\epsilon^{-2})$.

- (ii) *There exist $\Phi_1 := \begin{bmatrix} -i\phi_1 + \phi_2 \\ \phi_1 + i\phi_2 \end{bmatrix} + O(\epsilon^2)$, $\Phi_2 := \bar{\Phi}_1$ and purely imaginary numbers $\omega_1 := i(\lambda\epsilon^2 + O(\epsilon^3))$ and $\omega_2 := \bar{\omega}_1$ such that for $j = 1, 2$, the subspace $\mathbf{E}_j$ is spanned by Φ_j and*

$$\mathbf{H}_0\Phi_j = \omega_j\Phi_j, \quad \langle J\Phi_1, \Phi_1 \rangle = -4i, \quad \langle J\Phi_1, \Phi_2 \rangle = 0.$$

The projection $\mathbf{P}_j(\mathbf{H}_0) : \mathbf{E} \rightarrow \mathbf{E}_j$ is defined by

$$\mathbf{P}_1(\mathbf{H}_0)f = -\frac{1}{4i}\langle J\Phi_1, f \rangle \Phi_1, \quad \mathbf{P}_2(\mathbf{H}_0)f = \frac{1}{4i}\langle J\Phi_2, f \rangle \Phi_2, \quad \forall f \in \mathbf{E}.$$

- (iii) *There exist $\Phi_3 := \begin{bmatrix} \phi_0 \\ -i\phi_0 \end{bmatrix} + O(\epsilon^2)$, $\Phi_4 := \bar{\Phi}_3$ and purely imaginary numbers $\omega_3 = i(e_1 - e_0 + O(\epsilon^2))$, $\omega_4 = \bar{\omega}_3$ such that for $j = 3, 4$ the subspaces $\mathbf{E}_j$ is spanned by Φ_j and*

$$\mathbf{H}_0\Phi_j = \omega_j\Phi_j, \quad \langle J\Phi_3, \Phi_3 \rangle = 2i, \quad \langle J\Phi_3, \bar{\Phi}_4 \rangle = 0.$$

The projection $\mathbf{P}_j(\mathbf{H}_0) : \mathbf{E} \rightarrow \mathbf{E}_j$ is defined by

$$\mathbf{P}_3(\mathbf{H}_0)f = \frac{1}{2i}\langle J\Phi_3, f \rangle \Phi_3, \quad \mathbf{P}_4(\mathbf{H}_0)f = -\frac{1}{2i}\langle J\bar{\Phi}_4, f \rangle \bar{\Phi}_4, \quad \forall f \in \mathbf{E}.$$

- (iv) $\mathbf{E}_c = \{g \in \mathbf{E} : \langle Jf, g \rangle = 0, \quad \forall f \in \mathbf{E}_j, \quad \forall j = 0, 1, 2\}$. *Its corresponding projection is $\mathbf{P}_c(\mathbf{H}_0) = \text{Id} - \sum_{j=0}^4 \mathbf{P}_j(\mathbf{H}_0)$.*




Figure 4: Spectrum of $\mathbf{H}_0$ in the non-resonant case.

Corollary 4.16. For $r = (r_1, r_2, r_3) \in \mathbb{R}^3$, and for $j = 0, 1, 2, 3$, let $\Phi_{0j}^r := r * \Phi_{0j} = e^{-Jr_1} \Phi_{0j} \circ \Gamma_1(r_2, r_3)$. Similarly, for $j = 1, 2, 3, 4$, let $\Phi_j^r := e^{-Jr_1} \Phi_j \circ \Gamma_1(r_2, r_3)$. Also, let $\mathbf{E}_j^r$ and $\mathbf{E}_c^r$ be exactly as in Theorem 4.15, for $j = 0, 1, 2, 3, 4$. Then $\mathbf{E}_j^r$ are invariant under $\mathbf{H}_r$, $\mathbf{E} = \bigoplus_{j=0}^4 \mathbf{E}_j^r \oplus \mathbf{E}_c^r$, and the projections from $\mathbf{E}$ into these subspaces are defined exactly as those corresponding projections in Theorem 4.15. Moreover,

$$\mathbf{H}_\gamma \Phi_{00}^r = \Phi_{01}^r, \quad \mathbf{H}_\gamma \Phi_{0j}^r = 0, \quad \mathbf{H}_\gamma \Phi_k^r = \omega_k \Phi_k^r, \quad \forall j = 1, 2, 3, \quad k = 1, 2, 3, 4.$$

4.3 Resolvent Estimates and Decay Estimates

In this section, we shall study the resolvent $R(z) = (\vec{\mathcal{L}} - z)^{-1}$ with $\vec{\mathcal{L}} := \vec{\mathcal{L}}_Q$ for $Q = Q_E$ or $\tilde{Q}_E$. Set $e_{01} := e_1 - e_0$. In the resonant case, let ω denote the eigenvalue in the first quadrant. Recall that $\omega = i\kappa + \gamma$ with $\kappa = e_{01} + O(\epsilon^2)$ and $\gamma = \gamma_0 \epsilon^4 + O(\epsilon^5)$ for some constant $\gamma_0 > 0$. Also, recall that we can write

$$\vec{\mathcal{L}} = JH + W, \quad W = \begin{bmatrix} W_1 & W_2 \\ W_3 & W_4 \end{bmatrix} = O(\epsilon^2), \quad H = H_0 - E.$$

Lemma 4.17 (Resolvent Estimates). Let $R(z) := (\vec{\mathcal{L}} - z)^{-1}$ be the resolvent of $\vec{\mathcal{L}}$, $\mathbf{B} := \mathbf{B}(\mathbf{E}_s, \mathbf{E}_{-s})$ for some $s > 3$ where $\mathbf{E}_s$ is defined in (2.4). Then we have a constant $C > 0$ independent of ϵ such that for $\tau \geq |E|$,

$$\|R(i\tau \pm 0)\|_{\mathbf{B}} + \|R(-i\tau \pm 0)\|_{\mathbf{B}} \leq C[(1 + \tau)^{-1/2} + (|\tau - \kappa| + \epsilon^4)^{-1}]. \quad (4.45)$$

Moreover, for $k = 1, 2$, we also have,

$$\left\| R^{(k)}(i\tau \pm 0) \right\|_{\mathbf{B}} + \left\| R^{(k)}(-i\tau \pm 0) \right\|_{\mathbf{B}} \leq C[(1 + \tau)^{-(1+k)/2} + |\tau - \kappa| + \epsilon^4]^{-1}, \quad (4.46)$$

where $R^{(k)}$ is the k^{th} derivatives of R .

Proof. We prove the lemma for $\vec{\mathcal{L}} = \mathbf{H}_r$ in resonant case. For the other cases, the proof is simpler so we skip. Without loss of generality, we may assume $\vec{\mathcal{L}} = \mathbf{H}_0$. First of all, we study the resolvent operator $R(z)$ for $z \in \mathbb{C}$ near $\pm ie_{01}$. Assume that $|z - ie_{01}| < \epsilon$ or $|z + ie_{01}| < \epsilon$. For $\Phi \in \mathbf{E}_s, U \in \mathbf{E}$ such that $(\vec{\mathcal{L}} - z)U = \Phi$, we want to estimate $\|U\|_{\mathbf{E}_{-s}}$. We write $U = \Phi^* + \xi$, where $\Phi^* = a\Phi_0^+ + b\Phi_0^-$ with $a, b \in \mathbb{C}$ and $\xi \perp \Phi_0^\pm$. Then the equation $(\vec{\mathcal{L}} - z)U = \Phi$ is equivalent to

$$\begin{cases} a[i(e_0 - E) + \frac{1}{2}\langle \Phi_0^+, W\Phi_0^+ \rangle - z] + \frac{b}{2}\langle \Phi_0^+, W\Phi_0^- \rangle + \frac{1}{2}\langle \Phi_0^+, W\xi \rangle = \frac{1}{2}\langle \Phi_0^+, \Phi \rangle, \\ b[-i(e_0 - E) + \frac{1}{2}\langle \Phi_0^-, W\Phi_0^- \rangle - z] + \frac{a}{2}\langle \Phi_0^-, W\Phi_0^+ \rangle + \frac{1}{2}\langle \Phi_0^-, W\xi \rangle = \frac{1}{2}\langle \Phi_0^-, \Phi \rangle, \\ \xi = R_0(z)\mathbf{P}_c[\Phi - W(\Phi^* + \xi)] + \sum_{j=1}^3 \frac{\langle \Phi_j^\pm, \Phi - W(\Phi^* + \xi) \rangle}{2[\pm i(e_1 - E) - z]} \Phi_j^\pm. \end{cases} \quad (4.47)$$

Here $R_0(z) = (JH - z)^{-1}$ which is well-defined if $\text{Re}(z) \neq 0$. From (4.29), we get

$$\sum_{j=1}^3 \frac{\langle \Phi_j^\pm, \Phi - W(\Phi^* + \xi) \rangle}{2[\pm i(e_1 - E) - z]} \Phi_j^\pm = \sum_{j=1}^3 \frac{\langle \Phi_j^\pm, \Phi - W\xi \rangle}{2[\pm i(e_1 - E) - z]} \Phi_j^\pm. \quad (4.48)$$

From this and (4.47), we can write ξ as $\xi = \xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5$ where

$$\begin{aligned} \xi_1 &:= -aR_0(z)\mathbf{P}_c W\Phi_0^+ - bR_0(z)\mathbf{P}_c W\phi_0^-, \\ \xi_2 &:= -R_0(z)W\xi_1 - \frac{1}{2[\pm i(e_1 - E) - z]} \sum_{j=1}^1 \langle \Phi_j^\pm, W\xi_1 \rangle \Phi_j^\pm \\ \xi_3 &:= R_0(z)\mathbf{P}_c \Phi + \frac{1}{2[\pm i(e_1 - E) - z]} \sum_{j=1}^1 \langle \Phi_j^\pm, \Phi \rangle \Phi_j^\pm, \\ \xi_4 &:= -R_0(z)W\xi_3 - \frac{1}{2[\pm i(e_1 - E) - z]} \sum_{j=1}^1 \langle \Phi_j^\pm, W\xi_3 \rangle \Phi_j^\pm \\ \xi_5 &:= -R_0(z)W(\xi - \xi_1 - \xi_3) \\ &\quad - \frac{1}{2[\pm i(e_1 - E) - z]} \sum_{j=1}^1 \langle \Phi_j^\pm, W(\xi - \xi_1 - \xi_3) \rangle \Phi_j^\pm \end{aligned} \quad (4.49)$$

Now, for $z = i\tau + \delta$ with $|\tau - (e_0 - e_1)| \leq \epsilon$ and $0 < \delta \leq \epsilon^5$. Then, $|\pm i(e_1 - E) - z| = O(|e_1 - e_0|)$. So, from Lemma 4.3 and (4.49), we obtain

$$\begin{aligned} \|\xi_1\|_{\mathbf{E}_{-s}} &\leq C\epsilon^2[|a| + |b|], \quad \|\xi_2\|_{\mathbf{E}_{-s}} \leq C\epsilon^4[|a| + |b|], \\ \|\xi_3\|_{\mathbf{E}_{-s}} &\leq C\|\Phi\|_{\mathbf{E}_s}, \quad \|\xi_4\|_{\mathbf{E}_{-s}} \leq C\epsilon^2\|\Phi\|_{\mathbf{E}_s}. \end{aligned} \quad (4.50)$$

Similarly, we also have

$$\|\xi_5\|_{\mathbf{E}_{-s}} \leq C\epsilon^2[\|\xi_2\|_{\mathbf{E}_{-s}} + \|\xi_4\|_{\mathbf{E}_{-s}} + \|\xi_5\|_{\mathbf{E}_{-s}}].$$

So,

$$\|\xi_5\|_{\mathbf{E}_{-s}} \leq C\epsilon^4[\|\Phi\|_{\mathbf{E}_s} + \epsilon^2(|a| + |b|)]. \quad (4.51)$$

From (4.50) and (4.51), we obtain

$$\|\xi\|_{\mathbf{E}_{-s}} \leq C[\|\Phi\|_{\mathbf{E}_s} + \epsilon^2(|a| + |b|)]. \quad (4.52)$$

Recall that

$$\tau_0 = i(e_0 - E) + \frac{1}{2}\langle \Phi_0^+, W\Phi_0^+ \rangle = i[\kappa + O(\epsilon^2)], \quad \text{Re } \tau_0 = 0. \quad (4.53)$$

Let $M = (M_{ij})_{i,j=1}^2$ be the 2×2 matrix defined by

$$\begin{aligned} M_{11} &:= \tau_0 - \frac{1}{2} \langle \Phi_0^+, W R_0(z) \mathbf{P}_c W \Phi_0^+ \rangle - z, \\ M_{12} &:= \frac{1}{2} \langle \Phi_0^+, W \Phi_0^- \rangle - \frac{1}{2} \langle \Phi_0^+, W R_0(z) \mathbf{P}_c W \Phi_0^- \rangle, \\ M_{21} &:= \frac{1}{2} \langle \Phi_0^-, W \Phi_0^+ \rangle - \frac{1}{2} \langle \Phi_0^-, W R_0(z) \mathbf{P}_c W \Phi_0^+ \rangle, \\ M_{22} &:= \bar{\tau}_0 - \frac{1}{2} \langle \Phi_0^-, W R_0(z) \mathbf{P}_c W \Phi_0^- \rangle - z. \end{aligned} \quad (4.54)$$

Also, let

$$X := \frac{1}{2} \begin{bmatrix} \langle \Phi_0^+, \Phi \rangle - \langle \Phi_0^+, W(\xi_3 + \xi_4) \rangle \\ \langle \Phi_0^-, \Phi \rangle - \langle \Phi_0^-, W(\xi_3 + \xi_4) \rangle \end{bmatrix}, \quad B = \frac{1}{2} \begin{bmatrix} \langle \Phi_0^+, W(\xi_2 + \xi_5) \rangle \\ \langle \Phi_0^-, W(\xi_2 + \xi_5) \rangle \end{bmatrix}. \quad (4.55)$$

Then, the first two equations of (4.47) become

$$M \begin{bmatrix} a \\ b \end{bmatrix} + B = X. \quad (4.56)$$

From (4.52) and (4.55), we have

$$|X| \leq C \|\Phi\|_{\mathbf{E}_s}, \quad |B| \leq C \epsilon^6 [\|\Phi\|_{E_s} + (|a| + |b|)]. \quad (4.57)$$

On the other hand, from (4.35), (4.36) and (4.53), we have

$$\begin{aligned} M_{11} &= i(\kappa - \tau + O(\epsilon^2)) - i\lambda^2(\phi_0 Q^2, (H - iz)^{-1} \mathbf{P}_c \phi_0 Q^2) \\ &\quad - 2i\lambda^2(\phi_0 |Q|^2, (H + iz)^{-1} \mathbf{P}_c \phi_0 |Q|^2) - \delta, \\ M_{22} &= -i(\kappa + \tau + O(\epsilon^2)) - \delta - \frac{1}{2} \langle \Phi_0^-, W R_0(z) \mathbf{P}_c W \Phi_0^- \rangle. \end{aligned} \quad (4.58)$$

Because of $|z - i(e_0 - e_1)| \leq \epsilon$ and the Fermi Golden Rule, we get

$$\begin{aligned} \operatorname{Im} M_{11} &= (\kappa - \tau) + O(\epsilon^2) = O(\epsilon), \\ \operatorname{Re} M_{11} &= \lambda^2 \operatorname{Im}(\phi_0 Q^2, (H - iz)^{-1} \mathbf{P}_c \phi_0 Q^2) + O(\epsilon^5) \geq \frac{\lambda^2 \lambda_0 \epsilon^4}{2}, \\ \operatorname{Im} M_{22} &= -(\kappa + \tau) + O(\epsilon^2) = O(|e_0 - e_1|), \\ \operatorname{Re} M_{22} &= O(\epsilon^4). \end{aligned} \quad (4.59)$$

Note that from the symmetry property of $\tilde{Q}$, we get

$$(\phi_0^2, (\operatorname{Re} \tilde{Q})^2 - (\operatorname{Im} \tilde{Q})^2) = O(\epsilon^4), \quad (\phi_0^2, \operatorname{Re} \tilde{Q} \operatorname{Im} \tilde{Q}) = 0.$$

So, it follows that

$$\langle \Phi_0^+, W\Phi_0^- \rangle = 2i\lambda(\phi_0^2, \tilde{Q}^2), \quad \text{Re } c_1 = 0, \quad \text{Im } c_1 = O(\epsilon^4).$$

From this and (4.54), we get

$$M_{12} = O(\epsilon^4), \quad M_{21} = O(\epsilon^4). \quad (4.60)$$

From (4.59) and (4.60), we can find a constant $C_0 > 0$ depending on λ, λ_0 and $|e_0 - e_1|$ such that

$$|\det M| \geq C_0^{-1}[|\kappa - \tau| + \epsilon^4].$$

From this, (4.56), (4.57), we obtain

$$\begin{aligned} |a| + |b| &\leq |M^{-1}(X - B)| \leq C_0[|\kappa - \tau| + \epsilon^4]^{-1} \{ \|\Phi\|_{\mathbf{E}_s} + \epsilon^6(|a| + |b|) \} \\ &\leq C_0[|\kappa - \tau| + \epsilon^4]^{-1} \|\Phi\|_{\mathbf{E}_s} + C_0\epsilon^2(|a| + |b|). \end{aligned}$$

Thus, for sufficiently small ϵ such that $C_0\epsilon^2 < \frac{1}{2}$, we get

$$|a| + |b| \leq 2C_0[|\tau - \kappa| + \epsilon^4]^{-1} \|\Phi\|_{\mathbf{E}_s}. \quad (4.61)$$

From (4.52) and (4.61), we get $\|U\|_{\mathbf{E}_{-s}} \leq C[1 + |\tau - \kappa| + \epsilon^4]^{-1} \|\Phi\|_{\mathbf{E}_s}$. In other words, we have

$$\|R(i\tau + 0)\|_{(\mathbf{E}_s, \mathbf{E}_{-s})} \leq C[1 + (|\tau - \kappa| + \epsilon^4)^{-1}], \quad \forall \tau \in \mathbf{R} : |\tau - (e_0 - e_1)| \leq \epsilon.$$

Also, the estimates of $\|R(i\tau - 0)\|_{\mathbf{B}}, \|R(-i\tau \pm 0)\|_{\mathbf{B}}$ can be obtained in a similar way. So, we have proved (4.45) for τ near $\pm e_{01}$.

Next, for $z = i\tau + 0$ where $\tau > e_{01} + \epsilon$ or $|E| \leq \tau \leq e_{01} - \epsilon$, from [13, Theorem 9.2], we have $\|R_0(z)\|_{\mathbf{B}} \leq C(1 + \tau)^{-1/2}$. Then, by a similar argument as we just did, we also obtain (4.45).

Finally, to prove (4.46), we follow the argument in [23, Lemma 2.5]. Basically, we obtain (4.46) by an induction argument and by differentiating the relation $R(z)[1 + WR_0(z)] = R_0(z)$ and using the relations $(1 + WR_0)^{-1} = 1 - WR$, $(1 + R_0W)^{-1} = 1 - RW$. The proof of the lemma is then complete. $\square$

Lemma 4.18 (Decay Estimates). *For any of our excited states Q , let $\vec{\mathcal{L}} = \vec{\mathcal{L}}_Q$. Then we have*

- (i) *There exists a constant $C > 0$ independent of ϵ such that for all $\eta \in \mathbf{E}_c(\vec{\mathcal{L}}) \cap H^k$, we have*

$$C^{-1} \|\eta\|_{H^k} \leq \left\| e^{t\vec{\mathcal{L}}} \eta \right\|_{H^k} \leq C \|\eta\|_{H^k}, \quad \forall \eta \in \mathbf{E}_c(\vec{\mathcal{L}}) \cap H^k, \quad k = 1, 2.$$

(ii) For all $p \in [2, \infty]$, there is a constant $C = C(\epsilon, p)$ such that for all $\eta \in \mathbf{E}_c(\vec{\mathcal{L}})$, we have

$$\left\| e^{t\vec{\mathcal{L}}} \eta \right\|_{L^p} \leq C(\epsilon) |t|^{-3(1/2-1/p)} \|\eta\|_{L^{p'}}, \quad \text{where } \frac{1}{p} + \frac{1}{p'} = 1.$$

Proof. To prove (i), let's define the quadratic form: $\mathcal{Q}[\psi] = \langle \vec{\mathcal{L}}\psi, J\psi \rangle = \langle \mathbf{K}\psi, \psi \rangle$, $\psi \in \mathbf{E}$. Recall $\mathbf{K}$ is self-adjoint. Then, for all $\psi \in \mathbf{E}$, we have

$$\begin{aligned} \frac{\partial}{\partial t} \mathcal{Q}[e^{\vec{\mathcal{L}}t}\psi] &= \frac{\partial}{\partial t} \langle \mathbf{K}e^{J\mathbf{K}t}\psi, e^{J\mathbf{K}t}\psi \rangle \\ &= \langle \mathbf{K}J\mathbf{K}e^{J\mathbf{K}t}\psi, e^{J\mathbf{K}t}\psi \rangle + \langle \mathbf{K}e^{J\mathbf{K}t}\psi, J\mathbf{K}e^{J\mathbf{K}t}\psi \rangle = 0. \end{aligned}$$

Therefore, $\mathcal{Q}[e^{\vec{\mathcal{L}}t}\psi] = \mathcal{Q}[\psi]$, for all $t \geq 0$ and for all $\psi \in \mathbf{E}$.

Next, we claim that there exists $C > 0$ such that

$$C^{-1} \|\eta\|_{H^1}^2 \leq \mathcal{Q}[\eta] \leq C \|\eta\|_{H^1}^2, \quad \forall \eta \in \mathbf{E}_c(\vec{\mathcal{L}}) \cap H^1. \quad (4.62)$$

The upper-bound is immediate, so we just need the lower bound (the ‘‘spectral gap’’). Recall $\mathbf{K} = H - JW$, with $W = O(\varepsilon^2)$. In the non-resonant cases, the lower bound follows, by a simple perturbation-theoretic argument, from the corresponding spectral gap for the reference operator H . However, in the resonant cases, the subspaces $\mathbf{E}_c(\vec{\mathcal{L}})$ and $\mathbf{E}_c(JH)$ are not close, and the argument is more subtle – so let us assume now we are in the resonant case. Define the subspaces

$$S_0 := \text{span} \left\{ \begin{bmatrix} \phi_j \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ \phi_j \end{bmatrix} \mid j = 1, 2, 3 \right\}, \quad S_1 := \text{span} \{ \Phi_a, \Phi_b \}, \quad S := S_0 \oplus S_1,$$

where for $\vec{\mathcal{L}} = \mathbf{L}_0$, $(a, b) = (1, 3)$, and for $\vec{\mathcal{L}} = \mathbf{H}_0$, $(a, b) = (3, 4)$ (i.e. one eigenfunction from $\mathbf{E}_+$, one from $\mathbf{E}_-$). Notice S is 8-dimensional. For any $v \in S_0$,

$$\frac{\langle v, \mathbf{K}v \rangle}{\|v\|_{L^2}^2} = e_1 - E + O(\varepsilon^2) = O(\varepsilon^2).$$

Similarly, if $v \in S_0$, $w \in S_1$,

$$\langle v, \mathbf{K}w \rangle = \langle \mathbf{K}v, w \rangle = O(\varepsilon^2) \|v\|_{L^2} \|w\|_{L^2},$$

and

$$\langle v, w \rangle = O(\varepsilon^2) \|v\|_{L^2} \|w\|_{L^2}$$

since $\Phi_{a,b} = a_1 \begin{pmatrix} \phi_0 \\ \pm i\phi_0 \end{pmatrix} + h$, for an order-one constant a_1 , and with $\|h\|_{L^\infty} = O(\varepsilon^2)$. Now, for $\Phi = \Phi_a$ or Φ_b and $\omega = \omega_a$ or ω_b :

$$\begin{aligned} \bar{\omega} \langle \Phi, J\Phi \rangle &= \langle \vec{\mathcal{L}} \Phi, J\Phi \rangle = \langle \mathbf{K}\Phi, \Phi \rangle = \langle \Phi, \mathbf{K}\Phi \rangle \\ &= -\langle \Phi, J\vec{\mathcal{L}} \Phi \rangle = -\omega \langle \Phi, J\Phi \rangle \end{aligned}$$

and since $\bar{\omega} \neq -\omega$, $\langle \Phi, J\Phi \rangle = 0$, and so

$$\langle \Phi_a, \mathbf{K}\Phi_a \rangle = \langle \Phi_b, \mathbf{K}\Phi_b \rangle = 0.$$

Finally, notice

$$\begin{aligned} \bar{\omega}_a \langle \Phi_a, J\Phi_b \rangle &= \langle \vec{\mathcal{L}} \Phi_a, J\Phi_b \rangle = \langle \mathbf{K}\Phi_a, \Phi_b \rangle = \langle \Phi_a, \mathbf{K}\Phi_b \rangle \\ &= \langle J\Phi_a, \vec{\mathcal{L}} \Phi_b \rangle = \omega_b \langle J\Phi_a, \Phi_b \rangle = -\omega_b \langle \Phi_a, J\Phi_b \rangle \end{aligned}$$

and since $\bar{\omega}_a \neq -\omega_b$, we have $\langle \Phi_a, J\Phi_b \rangle = 0$ and so

$$\langle \Phi_a, \mathbf{K}\Phi_b \rangle = 0.$$

Combining all these facts, we conclude that

$$v \in S \implies \frac{\langle v, \mathbf{K}v \rangle}{\|v\|_{L^2}^2} = O(\varepsilon^2).$$

We claim now that:

$$J\eta \perp S_0 \oplus \mathbf{E}_+ \oplus \mathbf{E}_- \implies \langle \eta, \mathbf{K}\eta \rangle \geq \frac{1}{4}|e_1| \|\eta\|_{L^2}^2. \quad (4.63)$$

Indeed, if not, then for any v in the 9-dimensional subspace $S \oplus \langle \eta \rangle$, we would have $\langle v, \mathbf{K}v \rangle < \frac{1}{2}|e_1| \|v\|_{L^2}^2$. To see this, we are using $\langle v, \mathbf{K}\eta \rangle = O(\varepsilon^2) \|v\|_{L^2} \|\eta\|_{L^2}$ if $v \in S$, $J\eta \perp S$, which is easily checked, as well as $\|v + \eta\|_{L^2}^2 \sim \|v\|_{L^2}^2 + \|\eta\|_{L^2}^2$ for $v \in S$, $J\eta \perp S_0 \oplus \mathbf{E}_+ \oplus \mathbf{E}_-$, which, in turn, follows from the non-degeneracy of the matrix $\langle \Phi_j, J\Phi_k \rangle$, and a little linear algebra. The mini-max principle would then imply that $\mathbf{K}$ has at least 9 eigenvalues (counting multiplicity) below $\frac{1}{2}|e_1|$, while standard perturbation theory shows, in fact, that there are just 8.

Now since the Φ_{0j} are small L^2 perturbations of elements of S_0 , we get easily from (4.63)

$$\eta \in \mathbf{E}_c \implies \langle \eta, \mathbf{K}\eta \rangle \geq C^{-1} \|\eta\|_{L^2}^2.$$

Then it is straightforward to upgrade this lower bound to H^1 by using $\mathbf{K} = \delta(-\Delta + |E|) + (1 - \delta)\mathbf{K} + \delta R$, with R a bounded multiplication operator, for suitably small δ . This yields (4.62). From this and since $\mathcal{Q}[e^{\vec{\mathcal{L}}_t}\eta] = \mathcal{Q}[\eta]$, we get

$$\mathcal{Q}[e^{\vec{\mathcal{L}}_t}\eta] = \mathcal{Q}[\eta] \sim \|\eta\|_{H^1}^2,$$

which proves (i) for $k = 1$.

A straightforward consequence of the H^1 estimate just proved is that:

$$\|\eta\|_{H^3}^2 \sim \left\| \vec{\mathcal{L}}\eta \right\|_{H^1}^2 \sim \mathcal{Q}[\vec{\mathcal{L}}\eta], \quad \forall \eta \in \mathbf{E}_c \cap H^3.$$

Since $\mathcal{Q}[\vec{\mathcal{L}}\eta] = \mathcal{Q}[e^{t\vec{\mathcal{L}}}\vec{\mathcal{L}}\eta]$, it follows that $\|\eta\|_{H^3} \sim \left\| e^{t\vec{\mathcal{L}}}\eta \right\|_{H^3}$. Then, by interpolation, we obtain $\|\eta\|_{H^2} \sim \left\| e^{t\vec{\mathcal{L}}}\eta \right\|_{H^2}$, which is (i) for $k = 2$.

To prove (ii), we follow the argument in [23, Section 2.7]. Let $H_* = -\Delta - E$, $A := JV + W$. Then, we have $\vec{\mathcal{L}} = JH_* + A$. Note that H_* has no bound states and A is localized. Now, we define the wave operator $W_+ = \lim_{t \rightarrow \infty} e^{-t\vec{\mathcal{L}}} e^{tJH_*}$. Using Lemma 4.17 and the argument in [6], it follows that W_+ maps $\mathbf{E}$ onto $\mathbf{E}_c(\vec{\mathcal{L}})$. Moreover, W_+ and its inverse (restricted to $\mathbf{E}_c(\vec{\mathcal{L}})$) are bounded $L^p \rightarrow L^p$. Then, from the intertwining property, we have

$$e^{t\vec{\mathcal{L}}}\mathbf{P}_c = W_+ e^{tJH_*} (W_+)^* \mathbf{P}_c.$$

From this and the decay estimates of e^{tJH_*} , we obtain (ii). $\square$

5 Proof of Theorem 1.2

We shall divide the proof of Theorem 1.2 into two cases. The first case is for $Q = \tilde{Q}_E$, and the second one is for $Q = Q_E$. Each of these cases will be divided into two sub-cases depending on whether $e_0 < 2e_1$ (resonant case) or $e_0 > 2e_1$ (non-resonant case). We will give the details of the proof of Theorem 1.2 for the resonant case. In the non-resonant case, the proof is similar, and therefore we only sketch it.

We draw heavily on [23] in this section.

5.1 Stable Directions for the Excited State $\tilde{Q}_E$

5.1.1 The Resonant Case

For $r = (r_1, r_2, r_3) \in \mathbb{R}^3$, let $Z_r := (\Phi_{01}^r, \Phi_{02}^r, \Phi_{03}^r)$, $G_r := \Phi_{00}^r$ where $\Phi_{0j}^r, j = 0, 1, 2, 3$ are defined as in Corollary 4.14. Also, let $N_r := (\Phi_1^r, \Phi_2^r, \dots, \Phi_6^r) \in \mathbf{E}^6$.

We shall construct a solution ψ of the equation (1.1) of the form

$$\vec{\psi} = \begin{bmatrix} \operatorname{Re} \psi \\ \operatorname{Im} \psi \end{bmatrix} = e^{JEt} [\vec{Q}_{E,r(t)} + h],$$

where

$$h = \begin{bmatrix} h_1 \\ h_2 \end{bmatrix} = k + \eta, \quad k = \begin{bmatrix} k_1 \\ k_2 \end{bmatrix} := a(t)G_r + b(t) \cdot N_r, \quad \eta \in \mathbf{E}_c^r$$

with $a \in \mathbb{C}$, $b := (b_1, b_2, \dots, b_6) \in \mathbb{C}^6$ and

$$\vec{Q}_{E,r} = e^{Jr_1} \begin{bmatrix} \operatorname{Re} \tilde{Q}_E \circ \Gamma_1(r_2, r_3) \\ \operatorname{Im} \tilde{Q}_E \circ \Gamma_1(r_2, r_3) \end{bmatrix}.$$

Since (1.1) is equivalent to

$$\partial_t \vec{\psi} = (\Delta - V)J\vec{\psi} + \lambda|\vec{\psi}|^2 J\vec{\psi},$$

we get

$$\partial_t \vec{Q}_{E,r} + \partial_t h = \mathbf{H}_r h + F_1, \quad F_1 := \lambda[2\vec{Q}_{E,r} \cdot h + |h|^2]Jh + \lambda|h|^2 J\vec{Q}_{E,r}. \quad (5.1)$$

Now, let $\hat{k} := \nabla_r[a \cdot G_r + b \cdot N_r] = (\hat{k}_1, \hat{k}_2, \hat{k}_3)$, we get

$$\begin{aligned} \partial_t \vec{Q}_{E,r} &= \dot{r} \cdot Z_r, \quad \partial_t h = \dot{r} \cdot \hat{k} + \dot{a}G_r + \dot{b} \cdot N_r + \partial_t \eta, \\ \mathbf{H}_r h &= a\Phi_{01}^r + \sum_{j=1}^6 \omega_j b_j \Phi_j^r + \mathbf{H}_r \eta. \end{aligned}$$

Therefore, (5.16) becomes

$$\dot{r} \cdot (Z_r + \hat{k}) - a\Phi_{01}^r + \dot{a}G_r + \dot{b} \cdot N_r + \partial_t \eta = \sum_{j=1}^6 \omega_j b_j \Phi_j^r + \mathbf{H}_r \eta + F_1. \quad (5.2)$$

Taking the inner products of (5.2) with $J\Phi_{00}^r, J\Phi_{02}^r, J\Phi_{03}^r$, we obtain

$$\begin{aligned} \langle J\Phi_{00}, \Phi_{01} \rangle \dot{r}_1 + \sum_{j=1}^3 \dot{r}_j \langle J\Phi_{00}^r, \partial_{r_j} k \rangle &= \langle J\Phi_{00}^r, F_1 \rangle + a \langle J\Phi_{00}, \Phi_{01} \rangle, \\ \langle J\Phi_{03}, \Phi_{02} \rangle \dot{r}_2 + \sum_{j=1}^3 \dot{r}_j \langle J\Phi_{03}^r, \partial_{r_j} k \rangle &= \langle J\Phi_{03}^r, F_1 \rangle, \\ \langle J\Phi_{02}, \Phi_{03} \rangle \dot{r}_2 + \sum_{j=1}^3 \dot{r}_j \langle J\Phi_{02}^r, \partial_{r_j} k \rangle &= \langle J\Phi_{02}^r, F_1 \rangle. \end{aligned}$$

In other words, we can write $M\dot{r}^T = A$, where $M = M_0 + M_1$ with

$$M_0 := \begin{bmatrix} \langle J\Phi_{00}, \Phi_{01} \rangle & 0 & 0 \\ 0 & \langle J\Phi_{03}, \Phi_{02} \rangle & 0 \\ 0 & 0 & \langle J\Phi_{02}, \Phi_{03} \rangle \end{bmatrix}, \quad M_1 := \begin{bmatrix} \langle J\Phi_{00}^r, \hat{k} \rangle \\ \langle J\Phi_{03}^r, \hat{k} \rangle \\ \langle J\Phi_{02}^r, \hat{k} \rangle \end{bmatrix}, \quad (5.3)$$

and $A^T := (\langle J\Phi_{00}^r, F_1 \rangle + a\langle J\Phi_{00}, \Phi_{01} \rangle, \langle J\Phi_{03}^r, F_1 \rangle, \langle J\Phi_{02}^r, F_1 \rangle)$. We shall show that the matrix M is invertible and therefore, r satisfies $\dot{r} = [M^{-1}A]^T$. Now, let $F_2 := \dot{r} \cdot \hat{k} = [M^{-1}A]^T \cdot \hat{k}$ and $F = F_1 - F_2$. Taking the inner product of (5.2) with $J\Phi_{01}$ and $J\Phi_j^r$ for $j = 1, 2, \dots, 6$, we obtain

$$\dot{a} = \langle J\Phi_{01}, \Phi_{00} \rangle^{-1} \langle J\Phi_{01}^r, F \rangle, \quad \dot{b} = B.$$

Here $B = \omega + B'$ where $\omega = (\omega_1, \omega_2, \dots, \omega_6)$ and $B' = (B'_1, B'_2, \dots, B'_6) \in \mathbb{C}^6$ with

$$\begin{aligned} B'_j &= (J\Phi_j, \Phi_j)^{-1} \langle J\Phi_j^r, F \rangle, \quad j = 1, 2. \\ B'_3 &= (J\Phi_6, \Phi_3)^{-1} \langle J\Phi_6^r, F \rangle, \quad B'_6 = (J\Phi_3, \Phi_6)^{-1} \langle J\Phi_3^r, F \rangle, \\ B'_4 &= (J\Phi_5, \Phi_4)^{-1} \langle J\Phi_5^r, F \rangle, \quad B'_5 = (J\Phi_4, \Phi_5)^{-1} \langle J\Phi_4^r, F \rangle. \end{aligned} \quad (5.4)$$

Moreover, applying the projection $\mathbf{P}_c(\mathbf{H}_r)$ onto (5.2), we obtain the equation of η

$$\partial_t \eta = \mathbf{H}_r \eta + \mathbf{P}_c(\mathbf{H}_r) F. \quad (5.5)$$

Since $\mathbf{H}_r$ is time dependent, it's not easy to estimate η directly from (5.5). To overcome this difficulty, we shall use the following transformation: Let $\tilde{\eta} = \mathbf{P}_c(\mathbf{H}_0)\eta$. Then (5.5) becomes

$$\partial_t \tilde{\eta} = \mathbf{H}_0 \tilde{\eta} + \mathbf{P}_c(\mathbf{H}_0) NL, \quad NL := \tilde{F} + \mathbf{P}_c(\mathbf{H}_r) F, \quad (5.6)$$

where $\tilde{F} = (\mathbf{H}_r - \mathbf{H}_0)\eta$. Moreover, we have

$$\begin{aligned} \mathbf{P}_c(\mathbf{H}_r) - \mathbf{P}_c(\mathbf{H}_0) &= \\ &= \sum_{j=0}^2 [\mathbf{P}_j(\mathbf{H}_0) - \mathbf{P}_j(\mathbf{H}_r)] + [\mathbf{P}_+(\mathbf{H}_0) - \mathbf{P}_+(\mathbf{H}_r)] + [\mathbf{P}_-(\mathbf{H}_0) - \mathbf{P}_-(\mathbf{H}_r)]. \end{aligned} \quad (5.7)$$

From this, Lemma 4.41, we see that $\mathbf{P}_c(\mathbf{H}_r) - \mathbf{P}_c(\mathbf{H}_0) = O(|r|)$. Since $\eta = \tilde{\eta} + [\mathbf{P}_c(\mathbf{H}_r) - \mathbf{P}_c(\mathbf{H}_0)]\eta$, $|r|$ is sufficiently small and ϵ is fixed, we can solve η in term of $\tilde{\eta}$ by

$$\eta = \mathbf{U}_r \tilde{\eta}, \quad \mathbf{U}_r := \sum_{j=0}^{\infty} [\mathbf{P}_c(\mathbf{H}_r) - \mathbf{P}_c(\mathbf{H}_0)]^j. \quad (5.8)$$

Now, for given $\eta_\infty \in \mathbf{E}_c(\mathbf{H}_0)$, let $\tilde{\eta} = e^{\mathbf{H}_0 t} \eta_\infty + g$, where g satisfies the equation

$$\partial_t g = \mathbf{H}_0 g + \mathbf{P}_c(\mathbf{H}_0) N L,$$

and we want $g(t) \rightarrow 0$ in some sense as $t \rightarrow \infty$. In summary, we shall construct a solution ψ of (1.1) as

$$\vec{\psi} = e^{JEt} [\vec{Q}_{E,r(t)} + a \Phi_{00}^r + b \cdot \Phi_r + \mathbf{U}_r(\xi + g)],$$

where a, b, r satisfy the system of equations

$$\begin{cases} \dot{a} &= \langle J\Phi_{01}, \Phi_{00} \rangle^{-1} \langle J\Phi_{01}^r, F \rangle, \\ \dot{b} &= B, \quad \dot{r} = [M^{-1}A]^T, \\ \dot{g} &= \mathbf{H}_0 g + \mathbf{P}_c(\mathbf{H}_0) N L. \end{cases} \quad (5.9)$$

Now, for $\delta > 0$ and sufficiently small, let

$$\begin{aligned} \mathcal{X} := \{ & (a, r, b, g) : [0, \infty) \rightarrow \mathbb{C} \times \mathbb{R}^3 \times \mathbb{C}^6 \times (\mathbf{E}_c(\mathbf{H}_0) \cap H^2) : \\ & |a(t)|, |b(t)|, |r_j(t)| \leq \delta^{7/4} (1+t)^{-2}, \quad j = 2, 3; \quad |r_1(t)| \leq 2\delta^{7/4} (1+t)^{-1}, \\ & \|g(t)\|_{\mathbf{E} \cap H^2} \leq \delta^{7/4} (1+t)^{-3/2} \}. \end{aligned}$$

Then, we define the map $\Omega : \mathcal{X} \rightarrow \mathcal{X}$ with $\Omega(a, r, b, g) = (a^*, r^*, b^*, g^*)$ as

$$\begin{aligned} a^*(t) &= \int_{-\infty}^t [\langle J\Phi_{01}, \Phi_{00} \rangle^{-1} \langle J\Phi_{01}^r, F \rangle](s) ds, \\ r^*(t) &= \int_{-\infty}^t [M^{-1}A]^T(s) ds, \\ b_j^*(t) &= \int_{-\infty}^t e^{\omega_j(t-s)} B_j'(s) ds, \quad j = 1, 2, \dots, 4, \\ b_k^*(t) &= e^{\omega_k t} b_k(0) + \int_0^t e^{\omega_k(t-s)} B_k'(s) ds, \quad k = 5, 6, \\ g^*(t) &= \int_{-\infty}^t e^{\mathbf{H}_0(t-s)} \mathbf{P}_c N L(s) ds. \end{aligned}$$

Here $F = F_1 - F_2$ where F_1 is defined in (5.1) and $F_2 = (M^{-1}B)^T \cdot \hat{k} = (M^{-1}B)^T \cdot \nabla_r k$. Moreover, $\eta = \mathbf{U}_r[e^{\mathbf{H}_0 t} \eta_\infty + g]$. Note also that for $k = 5, 6$, we have $\text{Re } \omega_k < 0$. Therefore, the terms $e^{\omega_k t} b_k(0)$ in the equations of b_k^* exponentially decay and so $b_k(0)$ can be freely chosen.

Lemma 5.1. *The map Ω is well-defined and it is a contraction map if δ is sufficiently small and*

$$\|\xi_\infty\|_{H^2 \cap W^{2,1}} \leq \delta \quad \text{and} \quad |b_k(0)| \leq \delta^2/4, \quad \forall k = 5, 6.$$

Proof. Recall that $\eta := \xi + \mathbf{U}_r g$, $\xi := \mathbf{U}_r e^{\mathbf{H}_0 t} \eta_\infty$. Since $\|\xi_\infty\|_{H^2 \cap W^{2,1}} \leq \delta$, we get

$$\|\xi(t)\|_{H^2} \leq C(\epsilon)\delta, \quad \|\xi(t)\|_{W^{2,\infty}} \leq C(\epsilon)\delta|t|^{-3/2}$$

Therefore,

$$\||\xi|^2 J\xi\|_{H^2} \leq C(\epsilon)\delta^3(1+t)^{-3}.$$

Now, define $F_0 := 2(\vec{Q}_{E,r} \cdot \eta)J\eta + |\eta|^2 J\vec{Q}_{E,r} + |\eta|^2 J\eta$. We have $\|F_0\|_{H^2} \leq \delta^2(1+t)^{-3}$. From this and (5.1), we see that F_0 is the main term of F_1 . So, we obtain

$$\|F_1\|_{H^2} \leq C(\epsilon)\delta^2(1+t)^{-3}. \quad (5.10)$$

From (5.3), we have

$$\|M_1\| \leq C \left\| \hat{k} \right\|_{L^2} \leq C[|a| + |b|] \leq C\delta^{7/4}(1+t)^{-2}.$$

Therefore, for sufficiently small δ , $M = M_0 + M_1$ is invertible and $M^{-1} = (I_3 + M_0^{-1}M)M_0^{-1}$, where I_3 is the identity 3×3 matrix. From this and the explicit formula of A , we obtain

$$\begin{aligned} \|(M^{-1}A)_1\| &\leq \frac{3}{2}[|a| + \|M_0\| \|F_1\|_{L^2_{-s}}] \leq \frac{3}{2}|a| + C(\epsilon) \|F_1\|_{L^2_{-s}}, \\ \|(M^{-1}A)_k\| &\leq 2 \|M_0\| \|F_1\|_{L^2_{-s}} \leq C(\epsilon) \|F_1\|_{L^2_{-s}}, \quad k = 2, 3. \end{aligned} \quad (5.11)$$

Then, it follows from (5.10) and (5.11) that

$$\|F\|_{H^2} \leq \|F_1\|_{H^2} + \|F_2\|_{H^2} \leq C(\epsilon)\delta^2(1+t)^{-3}. \quad (5.12)$$

On the other hand, we also have

$$\|\tilde{F}\|_{H^2} = \|(\mathbf{H}_r - \mathbf{H}_0)\eta\|_{H^2} \leq C\epsilon|r| \|\eta\|_{L^\infty} \leq C(\epsilon)\delta^2(1+t)^{-5/2}.$$

From this and (5.10), we get

$$\|NL\|_{H^2} \leq C(\epsilon)\delta^2(1+t)^{-5/2}. \quad (5.13)$$

From Lemma 4.18 and (5.13), we obtain

$$\|g^*(t)\|_{H^2} \lesssim C(\epsilon)\delta^2 \int_\infty^t \|NL(s)\|_{H^2} ds \leq C(\epsilon)\delta^2(1+t)^{-3/2} \leq \delta^{7/4}(1+t)^{-3/2}.$$

On the other hand, from (5.4), (5.11) and (5.12) and by direct computation, we also obtain

$$|a^*|, |b^*|, |r_j^*| \leq \delta^{7/4}(1+t)^{-2}, \quad \forall j = 2, 3 \quad \text{and} \quad |r_1^*(t)| \leq 2\delta^{7/4}(1+t)^{-1}.$$

Hence, Ω maps $\mathcal{X}$ into $\mathcal{X}$. Also, it is easily checked that Ω is a contraction map if δ is sufficiently small. So, the lemma follows. $\square$

Now, to complete the proof of the Theorem 1.2 for $Q = \tilde{Q}_E$ in the resonant case, we shall prove that

$$\|\psi_{as}(t) - \psi(t)\|_{H^2} \leq C(\epsilon)(1+t)^{-3/2}. \quad (5.14)$$

where ψ, ψ_{as} are defined by

$$\begin{aligned} \psi &= e^{JEt} [\vec{Q}_{E,r(t)} + aG_r + b \cdot N_r + \mathbf{U}_r(\xi + g)] \\ \psi_{as}(t) &= e^{JEt} [\vec{Q}_{E,0} + \xi], \quad \xi := e^{\mathbf{H}_0 t} \eta_0. \end{aligned}$$

We have

$$\psi_{as}(t) - \psi(t) = e^{JEt} [\vec{Q}_{E,0} - \vec{Q}_{E,r} + aG_r + b \cdot N_r + (1 - \mathbf{U}_r)\xi + \mathbf{U}_r g].$$

From Lemma 4.12 and (5.7), we get

$$\begin{aligned} \left\| \vec{Q}_{E,0} - \vec{Q}_{E,r} + aG_r + b \cdot N_r \right\|_{H^2} &\leq C(\epsilon)[|r| + |a| + |b|] \\ &\leq C(\epsilon)\delta^{7/4}(1+t)^{-1}, \\ \|\mathbf{U}_r g\|_{H^2} &\leq C(\epsilon) \|g\|_{H^2} \leq C(\epsilon)(1+t)^{-3/2}. \end{aligned}$$

On the other hand, using (5.7) again, we obtain

$$\|(1 - \mathbf{U}_r)\xi\|_{H^2} \leq C(\epsilon)|r| \|\xi\|_{H^2} \leq C(\epsilon)\delta^{11/4}(1+t)^{-3/2}.$$

Therefore, we obtain (5.14).

5.1.2 The Non-Resonant Case

We shall construct a solution ψ of the equation (1.1) of the form

$$\vec{\psi} = \begin{bmatrix} \text{Re } \psi \\ \text{Im } \psi \end{bmatrix} = e^{JEt} [\vec{Q}_{E,r} + h],$$

where

$$h = \begin{bmatrix} h_1 \\ h_2 \end{bmatrix} = k + \eta, \quad k = \begin{bmatrix} k_1 \\ k_2 \end{bmatrix} := a(t)G_r + b(t) \cdot N_r, \quad \eta \in \mathbf{E}_c^r$$

with $a \in \mathbb{C}$, $b := (b_1, b_2, \dots, b_4) \in \mathbb{C}^4$ and

$$\vec{Q}_{E,r} = e^{Jr_1} \begin{bmatrix} \text{Re } \tilde{Q}_E \circ \Gamma_1(r_2, r_3) \\ \text{Im } \tilde{Q}_E \circ \Gamma_1(r_2, r_3) \end{bmatrix}.$$

Here, $G_r = \Phi_{00}^r$ and $N_r = (\Phi_1^r, \dots, \Phi_4^r)$. As in the previous section, we also obtain the equation of a, b, η as

$$\begin{cases} \dot{a} &= \langle J\Phi_{01}, \Phi_{00} \rangle^{-1} \langle J\Phi_{01}^r, F \rangle, \\ \dot{b} &= B, \quad \dot{r} = [M^{-1}A]^T, \\ \dot{g} &= \mathbf{H}_0 g + \mathbf{P}_c(\mathbf{H}_0)NL. \end{cases} \quad (5.15)$$

where $g = \mathbf{U}_r[e^{\mathbf{H}_0 t} \eta_\infty + \eta]$ and F, M, A, NL are defined exactly the same way. The map Ω is defined exactly the same except we don't have the equation for b_5^* and b_6^* .

5.2 Stable Directions for the Excited State Q_E

5.2.1 The Resonant Case

For $r = (r_1, r_2, r_3) \in \mathbb{R}^3$, let $Z_r := (\Phi_{01}^r, \Phi_{02}^r, \Phi_{03}^r), G_r := (\tilde{\Phi}_{01}^r, \tilde{\Phi}_{02}^r, \tilde{\Phi}_{03}^r) \in \mathbf{E}^3$ where $\Phi_{0j}^r, \tilde{\Phi}_{0j}^r, j = 1, 2, 3$ are defined as in Corollary 4.8. Also, let $N_r := (\Phi_1^r, \Phi_2^r, \Phi_3^r, \Phi_4^r) \in \mathbf{E}^4$. We shall construct a solution ψ of the equation (1.1) of the form

$$\vec{\psi} = \begin{bmatrix} \text{Re } \psi \\ \text{Im } \psi \end{bmatrix} = e^{JEt} [\vec{Q}_{E,r} + h], \quad \text{where } h := k + \eta, \quad k := a(t) \cdot G_r + b(t) \cdot N_r$$

with $a := (a_1, a_2, a_3) \in \mathbb{C}^3$, $b := (b_1, b_2, \dots, b_4) \in \mathbb{C}^4$, $\eta \in \mathbf{E}_c^r$ and

$$\vec{Q}_{E,r} = e^{Jr_1} \begin{bmatrix} Q_E \circ \Gamma_0(r_1, r_2) \\ 0 \end{bmatrix}.$$

Since (1.1) is equivalent to

$$\partial_t \vec{\psi} = (\Delta - V)J\vec{\psi} + \lambda |\vec{\psi}|^2 J\vec{\psi}.$$

So, we obtain

$$\partial_t \vec{Q}_{E,r} + \partial_t h = \mathbf{L}_r h + F_1, \quad F_1 := \lambda [2\vec{Q}_{E,r} \cdot h + |h|^2]Jh + \lambda |h|^2 J\vec{Q}_{E,r}. \quad (5.16)$$

Let $\hat{k} := \nabla_r[a \cdot G_r + b \cdot N_r]$, we have

$$\begin{aligned} \partial_t \vec{Q}_{E,r} &= \dot{r} \cdot Z_r, \\ \partial_t h &= \partial_t \eta + \dot{a} \cdot G_r + \dot{b} \cdot N_r + \dot{r} \cdot \hat{k}, \\ \mathbf{L}_r h &= \mathbf{L}_r \eta + a \cdot Z_r + \sum_{j=1}^4 \omega_j b_j \Phi_j^r. \end{aligned}$$

Therefore, the equation (5.16) becomes

$$(\dot{r} - a) \cdot Z_r + \dot{r} \cdot \hat{k} + \dot{a} \cdot G_r + \sum_{j=1}^4 (\dot{b}_j - \omega_j b_j) \Phi_j^r + \partial_t \eta = \mathbf{L}_r \eta + F_1. \quad (5.17)$$

Now, taking the inner product of (5.17) with $\tilde{\Phi}_{0j}^r, j = 1, 2, 3$, we obtain

$$(M_0 + M_1) \dot{r} = M_0 a + A,$$

where

$$M_0 := \begin{bmatrix} \langle J\tilde{\Phi}_{01}, \Phi_{01} \rangle & 0 & 0 \\ 0 & \langle J\tilde{\Phi}_{02}, \Phi_{02} \rangle & 0 \\ 0 & 0 & \langle J\tilde{\Phi}_{03}, \Phi_{03} \rangle \end{bmatrix}, \quad M_1 := \begin{bmatrix} \langle J\tilde{\Phi}_{01}^r, \hat{k} \rangle \\ \langle J\tilde{\Phi}_{02}^r, \hat{k} \rangle \\ \langle J\tilde{\Phi}_{03}^r, \hat{k} \rangle \end{bmatrix}, \quad (5.18)$$

and $A := (A_1, A_2, A_3)^T$, where $A_j = \langle J\tilde{\Phi}_{0j}^r, F_1 \rangle$. As we will see, the matrix $(M_0 + M_1)$ is invertible. Therefore, we obtain

$$\dot{r} = (1 + M_0^{-1} M_1)^{-1} a + (1 + M_0^{-1} M_1) M_0^{-1} A. \quad (5.19)$$

Now, for $\eta_\infty \in \mathbf{E}_c(\mathbf{L}_0)$, let g be such that $\eta = \mathbf{U}_r[e^{\mathbf{L}_0 t} \eta_\infty + g]$ where $\mathbf{U}_r$ is defined exactly the same way as in (5.8). Taking the inner products of (5.17) with $J\Phi_{0j}^r, \Phi_k^r$ we also get

$$\begin{cases} \dot{a}_j &= \langle J\Phi_{0j}, \tilde{\Phi}_{0j} \rangle^{-1} \langle J\Phi_{0j}^r, F \rangle, \\ \dot{b}_k &= \omega_k b_k + B_k, \\ \dot{g} &= \mathbf{L}_0 g + \mathbf{P}_c(\mathbf{L}_0) NL, \end{cases} \quad (5.20)$$

where $F = F_1 - F_2, F_2 := \dot{r} \cdot \hat{k} = [(1 + M_0^{-1} M_1)^{-1} a + (1 + M_0^{-1} M_1) M_0^{-1} A]^T \cdot \hat{k}$, $NL = (\mathbf{L}_r - \mathbf{L}_0) \eta + \mathbf{P}_c(\mathbf{L}_r) F$ and B satisfies $|B| \leq \|F\|_{L^2_s}$. Now, we define

$$\begin{aligned} \mathcal{Y} := \{ & (a, r, b, g) : [0, \infty) \rightarrow \mathbb{C}^3 \times \mathbb{R}^3 \times \mathbb{C}^4 \times (\mathbf{E}_c(\mathbf{L}_r) \cap H^2) : \\ & |a(t)|, |b(t)| \leq \delta^{7/4} (1+t)^{-2}, \quad |r(t)| \leq 2\delta^{7/4} (1+t)^{-1}, \\ & \|g(t)\|_{H^2} \leq \delta^{7/4} (1+t)^{-3/2} \}. \end{aligned}$$

Let $\Omega : \mathcal{Y} \rightarrow \mathcal{Y}$ be defined as $\Omega(a, r, b, \eta) = (a^*, r^*, b^*, g^*)$ where

$$\begin{aligned} a_j^*(t) &= \int_{-\infty}^t \langle J\Phi_{0j}, \tilde{\Phi}_{0j} \rangle^{-1} \langle J\Phi_{0j}^r, F \rangle(s) ds, \quad j = 1, 2, 3, \\ r^*(t) &= \int_{-\infty}^t [(1 + M_0^{-1}M_1)^{-1}a + (1 + M_0^{-1}M_1)M_0^{-1}A](s) ds, \\ b_k^*(t) &= \int_{-\infty}^t e^{\omega_k(t-s)} B_k(s) ds, \quad k = 1, 3, \\ b_l^*(t) &= e^{\omega_l t} b_l(0) + \int_0^t e^{\omega_l(t-s)} B_k(s) ds, \quad l = 2, 4, \\ g^*(t) &= \int_{-\infty}^t e^{\mathbf{L}_0(t-s)} \mathbf{P}_c(\mathbf{L}_0) N L(s) ds. \end{aligned}$$

Note that $\text{Re}(\omega_l) < 0$ for $l = 2, 4$. Therefore, the terms $e^{\omega_l t} b_l(0)$ decay exponentially and we only need to require $|b_l(0)| \leq \delta^2/4$. Then, as in Subsection 5.1.1, we can show that there exist ϵ_0 and $\delta_0(\epsilon)$ such that for $0 < \epsilon \leq \epsilon_0$ and $0 < \delta \leq \delta_0(\epsilon)$, the map Ω is well-defined and contraction. Therefore, we obtain the solution ψ of (1.1) of the form

$$\psi(t) = e^{JEt} [\vec{Q}_{E,r} + a \cdot G_r + b \cdot N_r + \mathbf{U}_r(e^{\mathbf{L}_0} \eta_\infty + g)]$$

and ψ satisfies the Theorem 1.2.

5.2.2 The Non-Resonant Case

The construction of the solution ψ is exactly the same as that of subsection 5.2.1. The only difference is that $b \in \mathbb{C}^2$, not $\mathbb{C}^4$ as in the Subsection 5.2.1. The equation of b^* becomes

$$b^*(t) = \int_{-\infty}^t B(s) ds, \quad B = (B_1, B_2).$$

The equations for a^*, r^* and g^* are the same as those in subsection 5.2.1. The proof of Theorem 1.2 is now complete.

Acknowledgments

We thank T.-P. Tsai and K. Nakanishi for their interest in this work. S.G. acknowledges the support of NSERC under Grant 251124-07

References

- [1] S. AGMON, *Spectral properties of Schrödinger operators and scattering theory*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 2 (1975), no. 2, pp. 151-218
- [2] SIMON L. ALTMANN, *Rotations, quaternions, and double groups*, Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1986
- [3] V.S. BUSLAEV AND G.S. PEREL'MAN, *Scattering for the nonlinear Schrödinger equations: states close to a solitary wave*, St. Petersburg Math J. 4 (1993), pp. 1111-1142.
- [4] V.S. BUSLAEV AND G.S. PEREL'MAN, *On the stability of solitary waves for nonlinear Schrödinger equations. Nonlinear evolution equations*, Amer. Math. Soc. Transl. Ser. 2, 164 (1995), pp. 75-98
- [5] V.S. BUSLAEV AND G.S. PEREL'MAN, *On asymptotic stability of solitary waves for nonlinear Schrödinger equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 20 (2003), no. 3, pp. 419-475.
- [6] S. CUCCAGNA, *Stabilization of solutions to nonlinear Schrödinger equations*, Comm. Pure Appl. Math., 54 (2001), no. 9, pp. 1110-1145.
- [7] S. CUCCAGNA, *On asymptotic stability of ground states of nonlinear Schrödinger equations*, Rev. Math. Phys., 15 (2003) no.8, pp. 877-903.
- [8] Z. GANG AND I. M. SIGAL, *Relaxation of solitons in nonlinear Schrödinger equations with potential*, Adv. Math., 216(2007), no. 2, pp. 443-490.
- [9] Z. GANG AND M.I. WEINSTEIN, *Dynamics of Nonlinear Schrödinger/Gross-Pitaevskii Equations; Mass Transfer in Systems with Solitons and Degenerate Neutral Modes*, Anal. PDE, Vol. 1 (2008), no. 3, pp. 267-322.
- [10] M. GOLUBITSKY, I. STEWART AND D. G. SHAEFFER, *Singularities and Groups in Bifurcation Theory*, Vol. II, Applied Math. Sciences, Springer-Verlag, 1988.
- [11] M. GOLUBITSKY AND I. STEWART, *The Symmetry Perspective*, Birkhäuser, 2000.
- [12] S. GUSTAFSON, K. NAKANISHI AND T.-P. TSAI, *Asymptotic stability and completeness in the energy space for nonlinear Schrödinger equations with small solitary waves*, Int. Math. Res. Not., 66(2004), pp. 3559-3584.

- [13] A. JENSEN AND T. KATO, *Spectral properties of Schrödinger operators and time-decay of the wave functions*, Duke Math. J., 46(1979), no. 3, pp. 583-611.
- [14] K. NAKANISHI, T.V. PHAN AND T.-P. TSAI, *Small Solutions of nonlinear Schrödinger equations near first excited states*, Preprint.
- [15] K.N. SRINIVASA RAO, *The Rotation and Lorentz Groups and their Representations for Physicists*, John Wiley & Sons, 1988.
- [16] H.A. ROSE AND M.I. WEINSTEIN, *On the Bound States of the Nonlinear Schrödinger Equation with a Linear Potential*, Physica D, 30(1988), pp. 207-218.
- [17] D.H. SATTERINGER, *Branching in the Presence of Symmetry*, CBMS-NSF Regional Conference Series in Applied Mathematics - Vol 40, Society for Industrial and Applied Mathematics, 1983.
- [18] A. SOFFER AND M.I. WEINSTEIN, *Multichannel Nonlinear Scattering Theory for Nonintegrable Equations I*, Com. Math. Phys., 133(1990), pp. 119-146.
- [19] A. SOFFER AND M.I. WEINSTEIN, *Multichannel Nonlinear Scattering Theory for Nonintegrable Equations II*, J. Differential Equations, 98(1992), pp. 376-390.
- [20] T.-P. TSAI, *Asymptotic dynamics of nonlinear Schrödinger equations with many bound states*, J. Differential Equations, 192(2003) pp. 225-282.
- [21] T.-P. TSAI AND H.-T. YAU, *Asymptotic dynamics of nonlinear Schrödinger equations: resonance dominated and dispersion dominated solutions*, Comm. Pure Appl. Math., 55(2002), pp. 153-216.
- [22] T.-P. TSAI AND H.-T. YAU, *Relaxation of excited states in nonlinear Schrödinger equations*, Int. Math. Res. Not., no.31(2002), pp. 1629-1673.
- [23] T.-P. TSAI AND H.-T. YAU, *Stable directions for excited states of nonlinear Schrödinger equations*, Comm. Partial Differential Equations, 27(2002), no. 11&12, pp. 2363-2402.
- [24] T.-P. TSAI AND H.-T. YAU, *Classification of asymptotic profiles for nonlinear Schrödinger equations with small initial data*, Adv. Theor. Math. Phys., 6(2002), no.1, pp. 107-139.
- [25] M.I. WEINSTEIN, *Lyapunov stability of ground states of nonlinear dispersive evolution equations*, Comm. Pure Appl. Math. , 39(1986), pp. 51-68.
- [26] K. YAJIMA, *The $W^{k,p}$ continuity of wave operators for Schrödinger operators*, J. Math. Soc. Japan, 47(1995), no. 3, pp. 551-581.

Stephen Gustafson, gustaf@math.ubc.ca
Department of Mathematics, University of British Columbia, Vancouver, BC
V6T 1Z2, Canada

Tuoc Van Phan, phan@math.ubc.ca
Department of Mathematics, University of British Columbia, Vancouver, BC
V6T 1Z2, Canada